



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

ADMIRE® PRO Systemic Protectant

EPA Reg. No. 264-827

For Foliar Uses on Potato, Tobacco, Fruiting Vegetables, Globe Artichokes, Herbs, Brassica (Cole) Leafy Vegetables, Leafy Green Vegetables, Legume Vegetables, Root, Tuberos and Corn Vegetables, Strawberry, Banana and Plantain, Bushberry, Caneberry, Citrus, Coffee, Grape, Hop, Pome Fruit, Pomegranate, Stone Fruit, Tree Nuts, Tropical Fruit, Christmas Trees, and Poplar/Cottonwood Trees.

Label Expires: 04-21-2014

Supplemental Label

ADMIRE® PRO Systemic Protectant

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read this label and the product package label before using this product. This Supplemental Label must be in possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for ADMIRE® PRO Systemic Protectant.

USE DIRECTIONS FOR SPECIFIC CROPS

POTATO – FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids Colorado potato beetle Flea beetles Leafhoppers Potato psyllid	1.3
Potato – Foliar Applications Apply ADMIRE PRO Systemic Protectant through properly calibrated ground and aerial application equipment.	
Potato – Foliar Application Restrictions Minimum interval between foliar applications: 7 days Maximum foliar applied ADMIRE PRO Systemic Protectant allowed per crop season: 5.6 fluid ounces/Acre (0.2 lb AI/Acre)	

TOBACCO – FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids	0.7 – 1.4
Flea beetles Japanese beetle	1.4
Tobacco – Foliar Applications Apply ADMIRE PRO Systemic Protectant through properly calibrated ground and aerial application equipment.	
Tobacco – Foliar Application Restrictions Pre-Harvest Interval (PHI): 14 days Minimum interval between foliar applications: 7 days Maximum ADMIRE PRO Systemic Protectant allowed per crop season: 7.8 fluid ounces/Acre (0.28 lb AI/Acre)	

FRUITING VEGETABLES – FOLIAR 1/

Crops of Crop Group 8 plus Okra including: Eggplant, Ground cherry, Okra, Pepper (including bell, chili, cooking, pimento and sweet) Tomato, Pepinos, Tomatillo

Pests Controlled	Rate fluid ounces/Acre
Aphids Colorado potato beetle Leafhoppers Whiteflies	1.3 – 2.2
Pepper weevil (Pepper only)	2.2
Fruiting Vegetables – Foliar Applications Apply ADMIRE PRO Systemic Protectant through properly calibrated ground and aerial application equipment. Thorough coverage with direct contact of the spray material to the target pests is required for optimum control. For pepper weevil, apply specified dosage of ADMIRE PRO Systemic Protectant by ground equipment only, timing applications prior to a damaging population becoming established. Good coverage of foliage and fruit is necessary for optimum control. Applications of ADMIRE PRO Systemic Protectant must be incorporated into a full-season program, where alternations of effective products from multiple classes of chemistry and different modes of action are utilized in a blocked or windowed approach. For additional information, please contact your Bayer CropScience representative, Extension specialist or Crop Advisor.	
Fruiting Vegetables – Foliar Application Restrictions Pre-Harvest Interval (PHI): 0 day Minimum interval between foliar applications: 5 days Maximum ADMIRE PRO Systemic Protectant allowed per crop season: 6.7 fluid ounces/Acre (0.24 lb AI/Acre)	
^{1/} Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	

GLOBE ARTICHOKE – FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers	1.4 – 3.5
Globe Artichoke – Foliar Applications Apply ADMIRE PRO Systemic Protectant through properly calibrated ground and aerial application equipment.	
Globe Artichoke – Foliar Application Restrictions Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 14 days Maximum ADMIRE PRO Systemic Protectant allowed per season: 14.0 fluid ounces/Acre (0.5 lb AI/Acre)	

HERBS – FOLIAR

Crops of Crop Subgroup 19A including: Angelica, Balm (lemon balm), Basil (fresh and dried), Borage, Bumet, Camomile, Catnip, Chervil (dried), Chinese chive, Chive, Clary, Coriander (cilantro or Chinese parsley leaves), Costmary, Culantro (leaf), Curry (leaf), Dillweed, Horehound, Hyssop, Lavender, Lemongrass, Lovage (leaf), Marigold, Marjoram, Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.

Pests Controlled	Rate fluid ounces/Acre
Aphids Flea beetles Leafhoppers Whiteflies	1.2
Herbs – Foliar Applications Apply ADMIRE PRO Systemic Protectant through properly calibrated ground and aerial application equipment.	
Herbs – Foliar Application Restrictions Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 5 days Maximum ADMIRE PRO Systemic Protectant allowed per season: 3.6 fluid ounces/Acre (0.13 lb AI/Acre)	
Herbs – Foliar Application Note Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, Bayer CropScience strongly recommends that only small areas or numbers of plants of each be treated and evaluated prior to commercial use.	

BRASSICA (COLE) LEAFY VEGETABLES ^{1/} – FOLIAR

Crops of Crop Group 5 including: Broccoli, Broccoli raab (*rapini*), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (*gai lon*) broccoli, Chinese (*bok choy*) cabbage, Chinese (*napa*) cabbage, Chinese mustard (*gai choy*) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens

Pests Controlled	Rate fluid ounces/Acre
Aphids Flea beetles Leafhoppers Whiteflies	1.3
Brassica (Cole) Leafy Vegetables – Foliar Application Restrictions Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 5 days Maximum ADMIRE PRO Systemic Protectant allowed per application: 6.7 fluid ounces/Acre (0.24 lb AI/Acre)	
^{1/} Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	

LEAFY GREEN VEGETABLES ^{1/} – FOLIAR

Crops Of Crop Subgroup 4A plus Watercress including: Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Roquette), Chervil, Chrysanthemum (edible leaved and garland), Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach)), Watercress (commercial production only, applications must not be made to native cress growing in streams or other bodies of water) Watercress (upland)

Pests Controlled	Rate fluid ounces/Acre
Aphids Flea beetles Leafhoppers Whiteflies	1.3
Leafy Green Vegetables (Crop Subgroup 4A) – Foliar Applications For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following the application. Applications must be made to fully leafed-up canopies, only.	
Leafy Green Vegetables (Crop Subgroup 4A) – Foliar Application Restrictions Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 5 days Maximum ADMIRE PRO Systemic Protectant allowed per application: 6.7 fluid ounces/Acre (0.24 lb AI/Acre)	
^{1/} Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	

LEGUME VEGETABLES 1/, Except Soybean (Dry) – FOLIAR

Crops of Crop Group 6 including: Edible Podded and Succulent Shelled Pea and Bean and Dried Shelled Pea and Bean

Bean (*Lupinus* spp., includes grain lupin, sweet lupin, white lupin, and white sweet lupin)

Bean (*Phaseolus* spp., includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)

Bean (*Vigna* spp., includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, yardlong bean)

Pea (*Pisum* spp., includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea)

Other Beans and Peas [Broad bean (fava), Chickpea (garbanzo bean), Guar, Jackbean, Lablab bean (hyacinth bean), Lentil, Pigeon pea, Soybean (immature seed), Sword bean]

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers Whiteflies	1.2
Legume Vegetables – Foliar Application Restrictions Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 7 days Maximum ADMIRE PRO Systemic Protectant allowed per crop season: 3.7 fluid ounces/Acre (0.13 lb AI/Acre)	
^{1/} Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	

ROOT, TUBEROUS and CORM VEGETABLES ^{1/} – FOLIAR

Crops of Crop Group 1 (except sugarbeet) plus Kava including: Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Beet (garden) ^{2/}, Burdock (edible) ^{2/}, Canna (edible, Queensland arrowroot), Carrot ^{2/}, Cassava (bitter & sweet) ^{2/}, Celeriac ^{2/}, Chayote (root), Chervil (turnip-rooted) ^{2/}, Chicory ^{2/}, Chufa, Dasheen (taro) ^{2/}, Ginger, Ginseng, Horseradish, Kava ^{2/}, ^{3/}, Leren, Parsley (turnip-rooted), Parsnip ^{2/}, Radish ^{2/}, Oriental radish (diakon) ^{2/}, Rutabaga ^{2/}, Salsify (black) ^{2/}, Salsify (oyster plant), Salsify (Spanish), Skirret, Sweetpotato ^{2/}, Tanier (cocoyam) ^{2/}, Tumeric, Turnip ^{2/}, Yam bean (jicama, manioc pea), Yam (true) ^{2/}.

(For recommended applications on Potato see Field Crops section)

Pests Controlled	Rate fluid ounces/Acre
Aphids Flea beetles Leafhoppers Whiteflies	1.2
Root, Tuberos and Corm Vegetables – Foliar Application Restrictions Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 5 days Maximum ADMIRE PRO Systemic Protectant allowed per crop season on Radish: 1.2 fluid ounces/Acre (0.044 lb AI/A); Maximum ADMIRE PRO Systemic Protectant allowed per crop season on all other crops: 3.7 fluid ounces/Acre (0.13 lb AI/A). Maximum ADMIRE PRO Systemic Protectant applications per crop season: 1 on Radish; 3 on other crops.	
^{1/} Not for use on crops grown for seed unless allowed by state-specific supplemental labeling. ^{2/} Tops or greens from these crops may be utilized for food or feed. ^{3/} Use not permitted in California unless otherwise directed by Supplemental Labeling.	

STRAWBERRY – FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids Spittlebugs Whiteflies	1.3
Strawberry – Foliar Application Restrictions Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 5 days Maximum ADMIRE PRO Systemic Protectant allowed per crop season: 3.9 fluid ounces/Acre (0.14 lb AI/A); Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.	

BANANA and PLANTAIN – FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers Thrips	2.8
Banana and Plantain – Foliar Applications Apply specified dosage of ADMIRE PRO Systemic Protectant as a broadcast or directed spray to infested area insuring thorough coverage. ADMIRE PRO Systemic Protectant may be applied through properly calibrated ground or aerial application equipment. Aerial application of ADMIRE PRO Systemic Protectant may result in slower activity and reduced control relative to results from ground application.	
Banana and Plantain – Foliar Application Restrictions Pre-Harvest Interval (PHI): 0 day Minimum interval between applications: 14 days Maximum ADMIRE PRO Systemic Protectant allowed per crop season: 14.0 fluid ounces/Acre (0.5 lb AI/A)	

BUSHBERRY – FOLIAR**Crops of Crop Subgroup 13B Including:** Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Ligonberry, Salal

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers/Sharpshooters	1.0 – 1.4
Blueberry maggot Japanese beetle (adults) Thrips (foliage feeding thrips only)	2.1 – 2.8
Bushberry – Foliar Application Restrictions Pre-Harvest Interval (PHI): 3 days Minimum interval between applications: 7 days Maximum ADMIRE PRO SYSTEMIC PROTECTANT allowed per season: 14.0 fluid ounces/Acre (0.5 lb AI/Acre) Maximum number of ADMIRE PRO SYSTEMIC PROTECTANT applications per crop season: 5 Minimum application volume (water): 20.0 GPA – ground; 5.0 GPA – aerial. Do not apply pre-bloom or during bloom or when bees are actively foraging.	

CANEBERRY – FOLIAR**Crops of Crop Subgroup 13A including:****Blackberry** (*Rubus eubatus*, including bingleberry, black satin berry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, Lavacaberry, Loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, and varieties and/or hybrids of these)**Raspberry** (black and red, *Rubus occidentalis*, *Rubus strigosus*, *Rubus idaeus*)

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers Thrips (foliage feeding thrips, only)	2.8
Caneberry – Foliar Application Restrictions Pre-Harvest Interval (PHI): 3 days Minimum interval between applications: 7 days Maximum ADMIRE PRO Systemic Protectant allowed per season: 8.4 fluid ounces/Acre (0.3 lb AI/Acre) Do not apply pre-bloom or during bloom or when bees are actively foraging.	

CITRUS (Field) – FOLIAR**Crops of Crop Group 10 Including:** Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Satsuma mandarin, White sapote (*Casimiroa* spp), and other cultivars and/or hybrids of these.

Pests Controlled	Rate fluid ounces/Acre
Aphids Asian citrus psyllid Blackfly Leafhoppers/Sharpshooters Leafminers Mealybugs Scales Whiteflies	3.5 – 7.0
Pest Suppressed	
Thrips (foliage feeding thrips only)	3.5 – 7.0
Citrus (Field) – Foliar Applications Scales – time applications to the crawler stage. Treat each generation.	
Citrus (Field) – Foliar Application Restrictions Pre-Harvest Interval (PHI): 0 day Minimum interval between applications: 10 days Maximum ADMIRE PRO Systemic Protectant allowed per crop season: 14.0 fluid ounces/Acre (0.5 lb AI/A) Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.	

COFFEE – FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers Leafminer	2.8
Pest Suppressed	
Scales	2.8
Coffee – Foliar Application Apply specified dosage of ADMIRE PRO Systemic Protectant as a broadcast or directed spray to infested area insuring thorough coverage. ADMIRE PRO Systemic Protectant may be applied through properly calibrated ground or aerial application equipment. Aerial application of ADMIRE PRO Systemic Protectant may result in slower activity and reduced control relative to results from ground application.	
Coffee – Foliar Application Restrictions Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 7 days Maximum ADMIRE PRO Systemic Protectant allowed per season: 14.0 fluid ounces/Acre (0.5 lb AI/Acre) Do not apply pre-bloom or during bloom or when bees are actively foraging.	

GRAPE – FOLIAR

Including: American bunch grape, Muscadine grape and Vinifera grape

Pests Controlled	Rate fluid ounces/Acre
Leafhoppers/Sharpshooters Mealybugs	1.0 – 1.4
Pest / Disease Suppressed	
Grapeleaf skeletonizer	1.3 – 1.4
Grape – Foliar Applications ADMIRE PRO Systemic Protectant may be applied by ground application, only.	
Grape – Foliar Application Restrictions Pre-Harvest Interval (PHI): 0 days Minimum interval between applications: 14 days Maximum ADMIRE PRO Systemic Protectant allowed per season: 2.8 fluid ounces/Acre (0.1 lb AI/Acre)	

HOP – FOLIAR

Pest Controlled	Rate fluid ounces/Acre
Aphids	2.8
Hop – Foliar Application Restrictions Pre-Harvest Interval (PHI): 28 days Minimum interval between applications: 21 days Maximum ADMIRE PRO Systemic Protectant allowed per season: 8.4 fluid ounces/Acre (0.3 lb AI/Acre)	

POME FRUIT – FOLIAR**Crops Of Crop Group 11 Including:** Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear), Quince

Pests Controlled	Rate fluid ounces/Acre
Leafhoppers	1.4 – 2.8
Aphids (except woolly apple aphid) Apple maggot Leafminers San Jose scale	2.8
FOR PEAR, ONLY Mealybugs Pear psylla	7.0
Pome Fruit – Foliar Application Applications targeting apple maggot should be combined with manufacturer's listed rate of a sticker.	
Pome Fruit – Foliar Application Restrictions Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 10 days Maximum ADMIRE PRO Systemic Protectant allowed per crop season: 14.0 fluid ounces/Acre (0.5 lb AI/A) Do not apply pre-bloom or during bloom or when bees are actively foraging.	

POMEGRANATE – FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers/Sharpshooters Whiteflies	2.8
Pest Suppressed	
Scales	2.8
Pomegranate – Foliar Application Restrictions Pre-Harvest Interval (PHI): 7 day Minimum interval between applications: 7 days Maximum ADMIRE PRO Systemic Protectant allowed per season: 8.4 fluid ounces/Acre (0.3 lb AI/Acre). Do not apply pre-bloom or during bloom or when bees are actively foraging.	

STONE FRUIT – FOLIAR

Crops Of Crop Group 12 Including: Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson and Japanese), Plumcot, Prune (fresh and dried)

Pests Controlled	Rate fluid ounces/Acre
Aphids Green June beetle Japanese beetle Leafhoppers/Sharpshooters Plant bugs Rose chafer San Jose scale	1.4 – 2.8
Cherry fruit fly	2.0 – 2.8
Pests Suppressed	
Plum curculio Stink bugs	2.8
Stone Fruit – Foliar Application Restrictions – Apricot, Nectarine, Peach: Pre-Harvest Interval (PHI): 0 day Minimum interval between applications: 7 days Maximum ADMIRE PRO Systemic Protectant allowed per crop season: 8.4 fluid ounces/Acre (0.3 lb AI/A) Minimum application volume (water): 50 GPA – ground application; 25 GPA – aerial application. Do not apply pre-bloom or during bloom or when bees are actively foraging. Stone Fruit – Foliar Application Restrictions – Cherries, Plums, Plumcot, Prune: Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 10 days Maximum ADMIRE PRO Systemic Protectant allowed per crop season: 14.0 fluid ounces/Acre (0.5 lb AI/A) Minimum application volume (water): 50 GPA – ground application; 25 GPA – aerial application Do not apply pre-bloom or during bloom or when bees are actively foraging.	

TREE NUTS – FOLIAR

Crops of Crop Group 14 except Almond including: Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut [black and English]

Pests Controlled	Rate fluid ounces/Acre
Aphids (except Black pecan aphid) Leafhoppers/Sharpshooters Phylloxera sp. (leaf infestations) Spittlebugs Whiteflies	1.2 – 2.4
Black pecan aphid Mealybugs San Jose scale	2.8
Tree Nuts – Foliar Applications Applications for control of San Jose scale should be timed according to crawler stage, treating each successive generation. Two applications on a 10 to 14-day interval may be required to achieve control.	
Tree Nuts – Foliar Application Restrictions Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 6 days Maximum ADMIRE PRO Systemic Protectant allowed per crop season: 10.0 fluid ounces/Acre (0.36 lb AI/A) Minimum application volume (water): 50 GPA – ground application, 25 GPA – aerial application Do not apply pre-bloom or during bloom or when bees are actively foraging.	

TROPICAL FRUIT – FOLIAR

Including: Acerola, Atemoya, Avocado, Birida, Black sapote, Canistel, Cherimoya, Custard apple, Feijoa, Jaboticaba, Guava, Llama, Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan, Rambutan, Sapodilla, Soursop, Spanish lime, Star apple, Starfruit, Sugar apple, Wax jambu

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers/Sharpshooters Mealybugs Thrips (foliage feeding thrips only) Whiteflies	2.8
Pest Suppressed	
Scales	2.8
Tropical Fruit – Foliar Application Restrictions Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 10 days Maximum ADMIRE PRO Systemic Protectant allowed per crop season: 14.0 fluid ounces/Acre (0.5 lb AI/A) Do not apply pre-bloom or during bloom or when bees are actively foraging.	

CHRISTMAS TREE – FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids Adelgids Sawflies	1.4 – 2.8
Christmas Tree – Foliar Applications Gall-forming adelgids – time applications to coincide with full bud-swell or first bud-break of earliest bud-breaking trees. Once galls form spraying will be ineffective.	
Christmas Tree – Foliar Application Restrictions Minimum interval between applications: 7 days Maximum ADMIRE PRO Systemic Protectant allowed per season: 14.0 fluid ounces/Acre (0.5 lb AI/Acre)	

POPLAR / COTTONWOOD – FOLIAR ^{1/}

(includes members of the genus *Populus* grown for pulp or timber)

Pests Controlled	Rate fluid ounces/Acre
Aphids Cottonwood leaf beetle	1.4 – 2.8
Pest Suppressed	
Poplar / Cottonwood – Foliar Application Restrictions Minimum interval between applications: 10 days Maximum ADMIRE PRO Systemic Protectant allowed per crop season: 14.0 fluid ounces/Acre (0.5 lb AI/A) Do not apply pre-bloom or during bloom or when bees are actively foraging.	
^{1/} Use not permitted in California unless otherwise directed by supplemental labeling.	

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
 For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.
 For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*