



LIQUID ORGANICS QUICK CAL

FORMULATED AS AN AID FOR HELPING PREVENT CORK SPOT, BITTER PIT, AND OTHER CALCIUM DEFICIENCY DISORDERS IN FRUIT TREES, FIELD AND VEGETABLE CROPS.

GUARANTEED ANALYSIS

Calcium (Ca).....12.0%
12.0% Water Soluble Calcium (Ca)

Derived from: Calcium Chloride.

This product does not contain more than 23% Chloride (Cl).

GENERAL INFORMATION

QUICK CAL is most effective when plants are growing rapidly and least effective when dormant or suffering from stress. On plants with waxy leaf surfaces, use a nonionic wetting agent at recommended rates. Do not use while temperatures are exceeding 90⁰ F. or when plants are under stress conditions. Determine compatibility with other fertilizers and/or pesticides by use of a jar test before tank mixing. Apply the product with non-corrosive spray equipment containing parts made of stainless steel, P.V.C., plastic, or fiberglass.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>.

PROPOSITION 65 WARNING: WARNING: This product may contain chemicals known to the State of California to cause cancer, birth defects or other reproductive harm.

NOTICE OF WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, NOR IS ANY REPRESENTATIVE OF SELLER AUTHORIZED TO MAKE ANY SUCH WARRANTY OR MODIFY THESE TERMS. This warranty does not extend to the storage, handling or use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such storage, handling or use. Seller shall not be responsible for incidental or consequential damages, if any, resulting from a breach of warranty.

Store above 32⁰ F

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose throat or skin. Avoid contact with eyes, skin and clothing. In case of skin contact, immediately flush with water. In case of eye contact, wash eye thoroughly with water for 15 minutes and get medical attention immediately. Do not contaminate feed or foodstuffs. Do not apply to any area not specified on label. Do not reuse empty container Wash thoroughly with water and detergent. Puncture, make unusable. Dispose of in a safe place.

DIRECTIONS FOR USE:

Pour recommended amount of QUICK CAL into partly filled spray tank. Add balance of water to fill tank. Keep agitated while filling and spraying. Flush all spray equipment after use.

AGRICULTURE CROPS

SOIL APPLICATION RECOMMENDATIONS

QUICK CAL can be mixed and applied with compatible liquid fertilizers or water . Apply 1 to 2 gallons of QUICK CAL per acre. Determine compatibility when mixing with other liquid fertilizers and/or pesticide before proceeding.

FORLIAR SPRAY RECOMMENDATIONS

Because of the nature of calcium deficiencies, QUICK CAL should be applied in multiple applications at 10-14 day intervals during the rapid growing and/or fruiting stage. For best results, spray when the crop is in an active growing state: after irrigation or natural rainfall. Spray early in the morning or late afternoon, Midday sprays may not be effective because of excessive moisture evaporation.

FIELD & VETGTABLE CROPS:

Apply 2 to 4 quarts per acre in at least 25 gallons of water. Application should be made early in the growing season. This treatment should be repeated if crops continue to show deficiencies.

ORCHARD CROPS:

Apply ½ to 1-1/2 gallons per acre in at least 100 gallons of water. Application can be made throughout the growing season and can be combined with cover sprays.

GRAPES:

Apply 2 to 4 quarts per acre in 50 to 100 gallons of water beginning at the “soft-berry” state and repeat at 10 to 14 day intervals during the growing season.

DRIP IRRIGATION

The required amount of this product should be applied over the growing season of the crop via multiple applications 1-2 weeks apart.

Consult your State Agricultural Experiment Station or State Extension Service Specialist for complete details on the spraying program best suited to your local conditions.

Manufactured By:
Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164-0589

Weight per gallon 11.3 lbs (5.13 kg)
Net Contents: 2.5 gallons (9.46 L)
Net Weight: 28 lbs 4 oz. (12.81k)

(Logo) Registered Material For Use in Organic Agriculture
Washington State Dept. of Agriculture