



Bayer CropScience LP  
 P.O. Box 12014  
 2 T.W. Alexander Drive  
 Research Triangle Park, North Carolina  
 27709  
 1-866-99BAYER (1-866-992-2937)  
<http://www.bayercropscienceus.com>

**AXIOM® DF Herbicide**

**For Use on Corn**

**EPA Reg. No. 264-766**

**EPA SLN. No. MN-000003**

**24(c) Supplemental Label**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MINNESOTA**

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

AXIOM USE RATES FOR EARLY POSTEMERGENCE APPLICATIONS AXIOM DF RATE (OUNCES PER ACRE) <sup>1,2,3</sup>		
Corn Leaf Stage at Spike to 3-leaf		
SOIL ORGANIC MATTER CONTENT	SOIL TEXTURAL GROUP <sup>4</sup>	
	MEDIUM	FINE
<0.5%	Do Not Use	Do Not Use
0.5 to 1%	10 to 15 oz	17 to 20 oz
1 to 1.5%	15 to 18 oz	
1.5 to 3%		
>3%	17 to 20 oz	

<sup>1</sup> Use the higher rate of AXIOM within the applicable rate range under any of the following conditions: heavy surface plant residues, heavy weed pressure and/or when soil organic matter is at the upper end of the range.  
<sup>2</sup> If soil pH is > 7.4, use the lower rate of the rate range shown.  
<sup>3</sup> Not recommended for use on peat or muck soils (soils with 20% or more organic matter.)  
<sup>4</sup> For more information, refer to the "Rate Selection/Soil Texture" section of the label.  
<sup>5</sup> Do not apply postemergence to field seedcorn inbreds and sweetcorn.

**BURNDOWN WEED CONTROL BY AXIOM**

## ANNUAL BROADLEAF CONTROL

Beggarweed, Florida	<i>Desmodium tortuosum</i>
Carpetweed	<i>Mollugo, verticillata</i>
Chickweed, common	<i>Stellaria media</i>
Cocklebur, common	<i>Xanthium strumarium</i>
Galinsoga	<i>Galinsoga parviflorum</i>
Lambsquarter, common <sup>1</sup>	<i>Chenopodium album</i>
Lettuce, prickly	<i>Lactuca serriola</i>
Marestail	<i>Coryza canadensis</i>
Mustard, blue	<i>Chorispora tenella</i>
Mustard, wild	<i>Sinapsis arvensis</i>
Pennycress, field	<i>Thlaspi arvense</i>
Pigweed, spp.	<i>Amaranthus spp.</i>
Pineappleweed	<i>Matricaria matriciodes</i>
Purselane, common	<i>Portulaca oleracea</i>
Sheperdspurse	<i>Capsella bursa-pastoris</i>
Sicklepod	<i>Cassia obtusifolia</i>
Thistle, Russian	<i>Salsola kali</i>
Velvetleaf	<i>Abutilon theophrastis</i>
Waterhemp, common <sup>1</sup>	<i>Amaranthus rudis</i>
Waterhemp, tall <sup>1</sup>	<i>Amaranthus tuberculatus</i>

<sup>1</sup> Axiom will not provide acceptable burndown control of triazine-resistant biotypes.

**SOIL-APPLIED  
WEED CONTROL/SUPPRESSION BY AXIOM\***

**ANNUAL GRASS WEEDS**

Barnyardgrass	<i>Echinochloa crusgalli</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>
Crabgrass, smooth	<i>Digitaria ischaemum</i>
Cupgrass, woolly*	<i>Erichloa villosa</i>
Foxtail, giant	<i>Sateria faberii</i>
Foxtail, green	<i>Sateria viridis</i>
Foxtail, yellow	<i>Setaria glauca</i>
Goosegrass,	<i>Eleusine indica</i>
Johnsongrass (seedling)	<i>Sorghum halepense</i>
Lovegrass, India	<i>Eragrostis spp.</i>
Millet, wild proso*	<i>Panicum milliaceum</i>
Nutsedge, yellow*	<i>Cyperus esculentus</i>
Panicum, browntop	<i>Panicum fasciculatum</i>
Panicum, Texas*	<i>Panicum texacunum</i>
Sanbur, field*	<i>Cenchrus incertus</i>
Shattercane*	<i>Sorghum bicolor</i>
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>
Witchgrass	<i>Panicum capillare</i>

**ANNUAL BROADLEAF WEEDS**

Beggarweed, Florida	<i>Desmodium tortuosum</i>
Carpetweed	<i>Mullugo verticillata</i>
Galinsoga	<i>Galinsoga, spp.</i>
Jimsonweed*	<i>Datura, stramonium</i>
Ladysthumb*	<i>Polygonum persicaria</i>
Lambsquarters, common <sup>1</sup>	<i>Chenopodium album</i>
Marestail*	<i>Conyza, canadensis</i>
Mustard spp. *	<i>Brassica, spp.</i>
Nightshade, eastern black*	<i>Solanum ptycanthum</i>
Pigweed spp. <sup>1</sup>	<i>Amaranthus spp.</i>
Purselane, common	<i>Portulaca oleracea</i>
Pusley, Florida	<i>Richardia scabara</i>
Ragweed, common*	<i>Ambrosia artemisiifolia</i>
Sida, prickly*	<i>Sida spinosa</i>
Smartweed, Pennsylvania*	<i>Polygonum pennsylvanicum</i>
Spurge, spotted	<i>Euphorbia maculata</i>
Sunflower, common*	<i>Helianthus annuus</i>
Thistle, Russian*	<i>Salsola kali</i>
Velvetleaf*	<i>Albutilon theophrasti</i>
Waterhemp, common <sup>1,2</sup>	<i>Amaranthus rudis</i>
Waterhemp, tall <sup>1,2</sup>	<i>Amaranthus tuberculatus</i>

\* Suppressed weeds may be stunted in growth and/or be reduced in population as compared to non-treated areas. Commercially acceptable control may require the application of an appropriate postemergence herbicide.

<sup>1</sup> Axiom may not provide acceptable control of triazine resistant biotypes.

<sup>2</sup> Due to extended germination intervals, full season control may not always be achieved, especially following preplant surface applications. To compliment control, Axiom should be used in tank-mixtures or sequentially with other herbicides that provide control of these weed species.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577  
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at [www.bayercropscienceus.com](http://www.bayercropscienceus.com)

Bayleton is a registered trademark of Bayer.

07/27/04