

Broad Spectrum Insecticide and Miticide

Ecotec®

**KEEP OUT
OF REACH OF
CHILDREN
CAUTION**

See enclosed booklet for complete
Precautionary Statements and
Directions For Use.

Active Ingredients	
Rosemary Oil	10.00%
Peppermint Oil	2.00%
Other Ingredients*	88.00%
Total	100.00%

* Wintergreen Oil, Lactic acid,
n-butyl ester, Vanillin and Lecithin.

FIFRA 25(b) Exempt Pesticide

This product has not been registered by the United States Environmental Protection Agency. Brandt Consolidated, Inc. represents that it is exempt from registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA).

OMRI®
Listed

NET CONTENTS: 2.5 GAL PER BOTTLE. PACKED 2 X 2.5 GAL



1026-01-100

Manufactured for:
Brandt Consolidated, Inc.
2935 South Koke Mill Road
Springfield, Illinois 62711 USA

BRANDT

KEEP OUT OF REACH OF CHILDREN CAUTION - PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID

IF IN EYES: Flush with water for at least 15 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a physician if irritation persists.

IF ON SKIN OR CLOTHING: Wash exposed area with plenty of soap and water for at least 15 minutes. Remove contaminated clothing. Get medical attention if irritation persists.

IF INHALED: Remove person to fresh air.

IF SWALLOWED: Rinse mouth out with water. Seek medical attention if feeling sick or nauseous.

Should an irritation persist, contact a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid contact with eyes, skin and clothing. Avoid breathing vapors or spray mist. May be harmful if swallowed. Wear long-sleeved shirt and long pants, shoes and socks, protective eyewear, and chemical resistant gloves made of neoprene, nitrile or natural rubber. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse. Keep all unprotected persons and pets out of treatment area until sprays have dried.

ENVIRONMENTAL HAZARDS

This product is not toxic to wildlife and possesses minimum risk to fish and aquatic organisms. Avoid direct contact to water. This product may be toxic to bees so avoid application when bees are actively foraging.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flames. Store only in original container.

GENERAL INFORMATION

ECOTEC is a broad spectrum, contact insecticide/miticide. It controls pests such as, but not limited to: aphids, thrips, whiteflies, mites, cabbage looper, and armyworm.

DIRECTIONS FOR USE

Use only according to label instructions. Read the entire label before using this product.

Only protected handlers may be in the application area. Do not apply this product in a way that will contact workers or other persons either directly or through drift.

Plant Safety

The use directions of this product are based on the results of product testing programs on a wide variety of plants. However, it is impossible to test this product on all species and cultivars. The phytotoxicity of ECOTEC has been assessed on a wide variety of common ornamental plants with no phytotoxicity. However, not all plant species and their varieties and cultivars have been tested with ECOTEC applied alone or with a possible tank mix combinations, sequential pesticide treatments, and adjuvants and surfactants. Local conditions also can influence plant tolerance and may not match those under which testing has been conducted. Therefore, before using ECOTEC, test the products on a sample of the plant and/or seedling. Do not apply on plants when the temperature is 90° F or above.

THE USE OF ADJUVANTS (SPREADERS AND/OR PENETRANTS) IS HIGHLY RECOMMENDED FOR IMPROVING PERFORMANCE.

ECOTEC can be applied through most standard or field type sprayers after dilution and mixing with water in accordance with label instructions. For best results, spray coverage should be uniform and complete. Always read and follow label directions for adjuvants.

Preharvest Interval

ECOTEC may be applied to crops at any time up to and including the day of harvest.

Use lower rates for light to moderate insect infestations and higher rates for heavy infestations. Repeat application as required to maintain the desired level of control.

Mixing and Application Instructions

SHAKE WELL BEFORE USING. TO MAINTAIN A HOMOGENEOUS SPRAY MIXTURE, CONSTANT AGITATION OR RECIRCULATION IS REQUIRED.

Apply spray solutions in properly maintained and calibrated equipment capable of delivering the desired volumes. Always clean tank, pump and lines thoroughly with water after use.

Mixing

Add ECOTEC to one-half (1/2) tank* under agitation containing water of 45° F or greater before filling to desired level. When combining with other products, such as adjuvants, insecticides, fungicides, or fertilizers, add these products first when the tank is approximately 1/2 full. Ensure that there is good agitation while mixing for complete emulsification. Maintain agitation during spray application.

* In areas with hard water (water with a total hardness of greater than 200 ppm) it is recommended to completely fill tank with water first and then add a water softener, mix thoroughly, add the appropriate quantity of ECOTEC, and mix thoroughly. It is critical that the water softener be added to the entire quantity of water and mixed thoroughly before introducing the ECOTEC to achieve maximum emulsion.

Application Equipment and Techniques

Spray droplet size should be adjusted to maximize coverage, minimize drift, and allow application to the intended exposure area only.

Boom Equipment

For best insect control using hydraulic or hydraulic air assist equipment, use the recommended rate of this product in sufficient gallons of spray solution per acre as a broadcast spray. Spray should thoroughly wet foliage but not run off.

Hand-held and High Volume Equipment

Spray-to-wet applications of this product may be made with knapsack and backpack sprayers, pump-up pressure sprayers, handguns, hand wands, and other hand-held spray equipment and vehicle mounted high volume spray equipment. For the most efficient use of spray mixture, spray all leaf surfaces uniformly and completely to wetness.

Aerial Applications

- Prior to application, check the aircraft/helicopter spray equipment for uniformity of spray pattern, spray width and output.
- Apply a minimum of 15 gallons per acre of diluted product.
- Optimum results are obtained when plant foliage is not too dense.
- **COVERAGE IS ESSENTIAL FOR GOOD CONTROL.**
- Use an output that will ensure optimum product penetration and coverage.
- Make aerial applications when the wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph.
- Low humidity and high temperature increase the evaporation rate of spray droplets and therefore the likelihood of good coverage.
- Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orientating nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 feet above the crop canopy is not recommended.

STORAGE AND DISPOSAL

Do not contaminate water, feed, or foodstuffs by storage or disposal.

Storage: Keep container tightly closed when not in use. Store only in the original container in a cool dry place away from children and pets. Keep from freezing.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on-site or at an appropriate waste disposal facility.

Container Disposal: Do not reuse empty container. Triple rinse empty container then puncture and dispose of container in a sanitary landfill, or incinerator, or if allowed by state and local regulations, by burning. If burned, stay out of smoke. Triple rinsed containers may be offered for recycling.

AGRICULTURE APPLICATION RATES

Spray when pests first appear. Repeat application every 5-7 days.

CROPS INCLUDING, BUT NOT LIMITED TO	INSECTS/MITES INCLUDING, BUT NOT LIMITED TO	APPLICATION RATE IN 100 GALLONS PER ACRE FOR GROUND APPLICATIONS
Legume & Fruiting Vegetables Beans, Eggplants, Lentils, Peas, Peanuts, Peppers, Tomatoes	Aphids, Whiteflies, Beetles, Two-Spotted Spider Mite, Plant Bugs, Thrips and Early Stages of Caterpillars	1.0 - 4.0 Pints
Bulb, Cole & Leafy Vegetables Broccoli, Cabbage, Cauliflower, Celery, Garlic, Lettuce, Onions, Spinach	Aphids, Whiteflies, Beetles, Two-Spotted Spider Mite, Plant Bugs, Thrips and Early Stages of Caterpillars	1.0 - 4.0 Pints For onion: Use 1.0 - 5.0 pints.
Cucurbits Cantaloupes, Cucumbers, Melons, Squashes, Watermelons	Aphids, Whiteflies, Beetles, Two-Spotted Spider Mite, Plant Bugs, Thrips and Early Stages of Caterpillars	1.0 - 4.0 Pints
Root & Tuber Vegetables Carrots, Potatoes, Radishes, Sweet Potatoes, Turnips	Aphids, Plant Bugs, Potato Leafhoppers, Two-Spotted Spider Mite and Thrips	1.0 - 4.0 Pints
Alfalfa	Aphids, Leafhoppers, Plant Bugs, Thrips and Early Stages of Caterpillars	1.5 - 4.0 Pints
Herbs & Spices Anise, Basil, Dill, Mint, Peppermint, Spearmint, Thyme, Wintergreen	Aphids, Leafhoppers, Plant Bugs, Thrips and Early Stages of Caterpillars	1.0 - 3.0 Pints
Small Fruits & Berries Blackberries, Blueberries, Cranberries, Raspberries, Strawberries	Aphids, Leafhoppers, Mealybugs, Whiteflies, Lygus Bugs, Thrips, Two-Spotted Spider Mite, Pacific Spider Mite, McDaniel Mite, European Red Mite and Early Stages of Leafrollers and Caterpillars	1.0 - 4.0 Pints
Grapes and Hops Grapes (Raisin, Table, Wine)	Leafhoppers, Sharpshooters, Two-Spotted Spider Mite, Pacific Spider Mite, Willamette Mite, Thrips, Mealybugs and Early Stages of Leafrollers and Caterpillars	1.0 - 4.0 Pints
Stone Fruits, Pome Fruits Apples, Apricots, Cherries, Peaches, Nectarines, Olives, Pears, Plums, Prunes, Pomegranates and Persimmons	Aphids, Thrips, Leafhoppers, Psylla, Scales, Two-Spotted Spider Mite, Pacific Mite, McDaniel Mite, European Red Mite, Early Stages of Leafrollers and Caterpillars	1.0 - 4.0 Pints
Citrus and Sub Tropicals Avocados, Lemons, Grapefruit, Oranges, Papayas, Bananas, Plantains, Mangos	Aphids, Thrips, Scales, Two-Spotted Spider Mite, Pacific Mite, Citrus Red Mite, Citrus Rust Mite, European Red Mite, Early Stages of Leafrollers and Caterpillars	2.0 - 5.0 Pints

For aerial applications use a minimum of 15 gallons of spray volume per acre. Complete coverage of foliage is essential to achieve control. The foliage should be completely WET, but avoid runoff. The use of a spreader and/or penetrant adjuvant improves product performance.

AGRICULTURE APPLICATION RATES

Spray when pests first appear. Repeat application every 5-7 days.

CROPS INCLUDING, BUT NOT LIMITED TO	INSECTS/MITES INCLUDING, BUT NOT LIMITED TO	APPLICATION RATE IN 100 GALLONS PER ACRE FOR GROUND APPLICATIONS
Nuts Almonds, Cashews, Pecans, Pistachios, Walnuts	Aphids, Thrips, Two-Spotted Spider Mite, Pacific Mite, McDaniel Mite and European Red Mite	1.0 - 4.0 Pints
Field Crops Corn, Cotton, Sorghum, Soybeans, Sweet Corn, Tobacco	Aphids, Soybean Aphid, Beetles, Whiteflies, Two-Spotted Spider Mite, Plant Bugs, Thrips, and Early Stages of Caterpillars	2.0 - 4.0 Pints
Christmas Trees Caution: Certain adjuvants may affect coloration of blue spruce	Mites and Aphids	1.0 - 4.0 Pints
Backyard Fruit and Nut Trees	Mites and Aphids	12 - 48 fl oz. Use sufficient water to get thorough coverage.
Mushroom Houses	Mushroom flies (Phorids and Sclarids)	4 - 8 fl oz per gallon - Spray dilution on outside and inside walls, floors, and sideboards of mushroom houses after compost has been pasteurized by heating. Also spray over the plastic covering the beds and trays after spawning. Retreat as needed.

For aerial applications use a minimum of 15 gallons of spray volume per acre. Complete coverage of foliage is essential to achieve control. The foliage should be completely WET, but avoid runoff. The use of a spreader and/or penetrant adjuvant improves product performance.

PESTS CONTROLLED THROUGH DRIP IRRIGATION SYSTEMS

(including but not limited to): Cutworms, Wireworms, Root Maggots, Symphylans, and Crickets.

CROPS TREATED BY DRIP IRRIGATION (including but not limited to): Beets, Brussel Sprouts, Cabbage, Chinese Broccoli, Broccoli, Chinese Cabbage, Chinese Mustard, Cauliflower, Collard, Endive, Garlic, Ginseng, Kale, Melons (Cantaloupes, Casabas, Honeydew, Muskmelons, Persians, and Watermelons), Onions (bulb and green), Parsnips, Potatoes, Sweet Potatoes, Spinach, Strawberries, Sugar Beets, Sweet Corn, Tomatoes.

Apply ECOTEC through the drip irrigation systems at a rate of 2 pints per acre of plant bed as a preventative treatment or when pest infestation is low to moderate and 4 pints per acre of plant bed when pest infestation is high. Apply 2 to 3 applications of ECOTEC over a 7 to 14 day schedule. Refer to local extension recommendations for additional information regarding pest populations and thresholds.

ECOTEC can be applied at any time during the crop cycle, from the time of planting to the harvest and there are no restrictions on the number of applications per season. For drip irrigation systems, set the emitter spacing and the line pressure to give the best bed coverage and reduce loss of chemigation products through leaching.

BED SPACING	LINEAR FEET OF BED TO EQUAL ONE ACRE	ECOTEC, 2 PINTS/ACRE RATE/1000 ROW FEET	ECOTEC, 4 PINTS/ACRE RATE/1000 ROW FEET
36 inches	14,520 ft	2.2 fl oz	4.4 fl oz
48 inches	10,890 ft	2.9 fl oz	4.4 fl oz
60 inches	8,712 ft	3.7 fl oz	4.4 fl oz
72 inches	7,260 ft	4.4 fl oz	4.4 fl oz

USE RATES FOR ANY PLANTS GROWN INDOORS OR IN GREENHOUSES, SHADEHOUSES, INTERIORSCAPES, AND NURSERIES

Spray when pests first appear. Repeat application every 5-7 days.

PESTS INCLUDING, BUT NOT LIMITED TO	FL OZ/10 GAL SPRAY SOLUTION	FL OZ/100 GAL SPRAY SOLUTION
APHIDS: Green Peach aphids, Pea aphids, Cotton aphids, Rose aphids	2 - 4	20 - 40
BEETLES: Flea beetles, Leaf beetles	2.0 - 4.8	20 - 48
Early Stages of Caterpillars: Armyworms, Leafrollers, Loopers, Webworms	1.6 - 3.2	16 - 32
FLIES: Fungus gnats, Shore flies	1.6 - 3.2	16 - 32
LEAFHOPPERS: Grape leafhoppers, Potato leafhoppers	1.6 - 3.2	16 - 32
LEAF MINERS: Azaela leaf miners, Citrus leaf miners, Blotch leaf miners, Boxwood leaf miners	1.6 - 3.2	16 - 48
MEALYBUGS: Citrus mealybugs, Long-tailed mealybugs	1.6 - 3.2	16 - 48
MITES: Two-Spotted Spider mite, Broad mite, Citrus budmite, Cyclamen mite, Rust mite	1.6 - 4.8	16 - 48
SOFT SCALES (Crawler Stages)	1.6 - 4.8	16 - 48
THRIPS: Western flower thrips	3.2 - 6.4	32 - 64
WHITEFLIES: Greenhouse whiteflies, Silverleaf whiteflies	1.6 - 3.2	16 - 32

For aerial applications use a minimum of 15 gallons of spray volume per acre. Complete coverage of foliage is essential to achieve control. The foliage should be completely WET, but avoid runoff. The use of a spreader and/or penetrant adjuvant improves product performance.

USE RATES FOR OUTDOOR PLANTS INCLUDING TREES, LANDSCAPES, TURFGRASS, NURSERY, AND ALL OUTDOOR ORNAMENTAL PLANTS

Spray when pests first appear. Repeat application every 5-7 days.

PEST INCLUDING, BUT NOT LIMITED TO	FL OZ/10 GAL SPRAY SOLUTION	FL OZ/100 GAL SPRAY SOLUTION
APHIDS: Green Peach aphids, Pea aphids, Cotton aphids, Rose aphids	2.0 - 4.0	20 - 40
BETTERLES: Colorado potato Flea beetles, Japanese beetles, Leaf beetles, Mexican bean beetles, Rose chafers, Twig girdlers	3.2 - 6.4	32 - 64
BUGS: Boxelder bugs, Chinch bugs, Lygus bugs, Spittle bugs	1.6 - 4.8	16 - 48
Early Stages of Caterpillars: Armyworms, Cutworms, Fruitworms, Gypsy moths, Hornworms, Leafrollers, Navel orange worms, Pine tip moths, Sod webworms, Tent caterpillars, Tobacco budworms, Tussock moths	1.6 - 3.2	16 - 32
FLIES: Cherry maggots, Crane flies, Midges	2.0 - 4.0	16 - 48
LEAFHOPPERS: Variegated leafhoppers and Sharpshooters	3.2 - 6.4	16 - 48
LEAF MINERS: Azaela leaf miners, Citrus leaf miners, Blotch leaf miners, Boxwood leaf miners	1.6 - 4.8	16 - 48
MEALYBUGS: Apple mealybugs, Citrus meal bugs, Grape mealybugs	1.6 - 4.8	16 - 48
MITES: Two-Spotted Spider mite, Broad mite, Citrus budmite, Cyclamen mite, Rust mite	1.6 - 4.8	16 - 48
PSYLLIDS: Pear psylla	1.6 - 4.8	16 - 48
SAWFLIES: European pine sawflies, Pear sawflies, Red-headed pine sawflies, Yellow-headed pine sawflies.	1.6 - 3.2	16 - 32
Crawler stages of SCALES: Black scales, Brown soft scales, California red scales, Coffee scales, Olive scales, San Jose scales	1.6 - 4.8	16 - 48
THRIPS: Citrus thrips, Onion thrips, Western flower thrips	3.2 - 6.4	32 - 64
WHITEFLIES: Greenhouse whiteflies, Silverleaf whiteflies, Woolly whiteflies	1.6 - 3.2	16 - 32

For aerial applications use a minimum of 15 gallons of spray volume per acre. Complete coverage of foliage is essential to achieve control. The foliage should be completely WET, but avoid runoff. The use of a spreader and/or penetrant adjuvant improves product performance.

Warranty

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use; but neither this warranty nor any other warranty of merchantability or fitness of a particular product expressed or implied, extends to the use of this product contrary to label conditions, or under conditions not reasonably foreseeable to the seller; and buyer assumes the risk of any such use.