

# Manzate<sup>®</sup> Pro-Stick<sup>™</sup> fungicide

## DISPERSIBLE GRANULES

### ACTIVE INGREDIENTS BY WEIGHT

A coordination product of zinc ion and manganese ethylenebisdithiocarbamate . . . . .	75.0%
in which the ingredients are:	
Manganese++ . . . . .	15.0%
Zinc++ . . . . .	1.9%
Ethylenebisdithiocarbamate ion (C <sub>4</sub> H <sub>6</sub> N <sub>2</sub> S <sub>4</sub> )-- . . . . .	58.1%

**INERT INGREDIENTS** . . . . . 25.0%

**TOTAL** . . . . . 100.0%

Contains 0.75 Pound of Mancozeb Per Pound of Product

EPA Reg. No. 70506-234

EPA Est. No. 352-COL-001

### KEEP OUT OF REACH OF CHILDREN CAUTION

#### FIRST AID

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **Contact the Rocky Mountain Poison Center at 1-866-673-6671 for emergency medical treatment information.** See Label for Additional Precautions and Directions for Use.

**FOR CHEMICAL EMERGENCY:**  
Spill, leak, fire, exposure, or accident,  
call CHEMTREC at 1-800-424-9300.

See inside for additional Precautionary Statements and complete Directions for Use.

Net Contents: 6 lbs., 30 lbs., 50 lbs.



United Phosphorus, Inc.  
630 Freedom Business Center, Suite 402  
King of Prussia, PA 19406  
1-800-438-6071 • www.upi-usa.com

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

MAY IRRITATE EYES, NOSE, THROAT AND SKIN. MAY BE HARMFUL IF ABSORBED THROUGH SKIN, INHALED OR SWALLOWED. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Keep away from fire or sparks.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

#### Applicators and other handlers (other than mixers and loaders) must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical resistant gloves, such as nitrile rubber, natural rubber, or butyl rubber
- Shoes plus socks

#### Mixers and Loaders must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical resistant gloves, such as nitrile rubber, natural rubber, or butyl rubber
- Shoes plus socks
- Protective eyewear
- Chemical-resistant apron when mixing or loading

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### ENGINEERING CONTROL STATEMENTS:

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

During aerial application, human flaggers must be in enclosed cabs.

#### USER SAFETY RECOMMENDATIONS

**USERS SHOULD:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or incorporate spilled treated seed. Do not contaminate water by disposing of equipment washwaters.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard

and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Chemical resistant gloves, such as nitrile rubber, natural rubber, or butyl rubber
- Shoes plus socks

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Commercial seed treatments are not within the scope of the Worker Protection Standard.

Do not enter treated areas until sprays have dried.

United Phosphorus, Inc. will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by United Phosphorus, Inc. User assumes all risks associated with such nonrecommended use.

MANZATE® PRO-STICK™, a dispersible granule containing mancozeb, is recommended for use as a spray for the control of many important plant diseases.

## APPLICATION INSTRUCTIONS

**AS A SPRAY** (Ground or Aerial Equipment) - Apply MANZATE® PRO-STICK™ at the rate shown; use sufficient water to provide thorough coverage: use 20 to 100 gallons per acre for ground equipment and no less than 2 gallons per acre for aircraft. Add MANZATE® PRO-STICK™ slowly to water in the spray tank with agitation, or premix thoroughly in separate holding tank for concentrate or aircraft sprayers. Continuous agitation is required to keep the product in suspension. A spreader-sticker spray adjuvant may be used with this product if needed; contact your local product distributor or United Phosphorus, Inc. representative for specific recommendations.

## RESTRICTIONS

### **Foliar Applications**

#### **Where EBDC Products Used Allow the Same Maximum Poundage of Active Ingredient Per Acre Per Season**

If more than one product containing an EBDC active ingredient (maneb, mancozeb or metiram) is used on a crop during the same growing season and the EBDC products used allow the same maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed any one of the specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

#### **Where EBDC Products Used Allow Different Maximum Poundage of Active Ingredient Per Acre Per Season**

If more than one product containing an EBDC active ingredient is used on a crop during the same growing season and the EBDC products used allow different maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed the lowest specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

### **SEED TREATMENT**

In addition to the maximum number of foliar applications permitted by the formula stated above, a single application for seed treatment may be made on crops which have registered seed treatment uses.

### **CHEMIGATION**

Apply MANZATE® PRO-STICK™ fungicide only through sprinkler systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation systems. Do not apply MANZATE® PRO-STICK™ through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

### **Specific Instructions for Public Water Systems:**

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

### **Specific Instructions for Sprinkler Irrigation Systems:**

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Good agitation is required in the injection tank.
9. In moving systems, apply specified dosage of MANZATE® PRO-STICK™ as a continuous injection. In non-moving systems inject MANZATE® PRO-STICK™ for 15 to 30 minutes at end of cycle. Use the least amount of water possible consistent with uniform coverage.
10. Mix the amount of MANZATE® PRO-STICK™ needed for acreage to be treated into the quantity of water determined during prior calibration. For moving systems inject into the system continuously for one complete revolution of the field. For non-moving systems inject into system for the time established during calibration.
11. Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all MANZATE® PRO-STICK™ is flushed from system.

<b>CROP</b>	<b>DISEASES CONTROLLED</b>	<b>RATE OF MANZATE® PRO-STICK™ PER APPLICATION LBS/ACRE</b>	<b>TIMING/INTERVALS (Also refer to Directions for Use)</b>	<b>RESTRICTIONS/COMMENTS</b>
Apple	(See Pomefruit)			
Asparagus	Cercospora Leaf Spot, Rust	2	Start applications when rust first appears and repeat at 10 day intervals. Four applications are usually sufficient.	Apply only on asparagus ferns after spears have been harvested. Do not apply within 180 days of harvest in all states except CA and AZ (120 days). Do not apply more than 8 lbs (6 lbs active) per acre per season.
Asparagus Crown (Planting Stock)	Crown Rot	1.0 lb/100 gals	Dip clean, loosely packed crowns into continuously agitated fungicide suspension for 5 minutes. Drain and plant as soon as possible.	Wash dirty crowns before dip treatment. Replace suspension in clean tank when discolored by soil.
Banana (Including Plantain)	Sigatoka	2-3	Apply when leaves first appear and repeat every 14 to 21 days or as required. Use sufficient water to provide adequate coverage.	Do not apply more than 30 lbs (22.5 lbs active) per acre per growing cycle. Minimum preharvest interval 0 days.
Barley, Oat, Rye, Wheat (Including Triticale)	Helminthosporium Leaf Spot, Leaf Rust, Septoria Glume Blotch, Septoria Leaf Spot, Tan Spot	2	Start application at onset of disease or when plants are in the tillering to jointing stage and repeat at 7 to 10 day intervals.	Do not make more than three applications during the season. Do not apply more than 6 lbs (4.5 lbs active) per acre per crop. Do not apply within 26 days of harvest. Do not graze livestock in treated areas prior to harvest.
Caprifig (Non-Food Use)	Endosepsis (Fusarium), Mold	4 lbs/100 gals	Prepare mamme figs by making a shallow cut through the eye and then hand dividing to avoid wasp injury. Submerge mamme figs in the continuously agitated suspension for at least 15 minutes. Drain before placement in trees.	Use fresh dipping suspension after treating 4 to 5 batches of figs.
Corn (Sweet Corn for Fresh Use or Processing; Popcorn; and Sweet Corn for Seed Production, including Hybrid Seed)	Common Rust, Helminthosporium Leaf Blight, Gray Leaf Spot	1.5	Use sufficient water for thorough coverage. Start applications when disease first appears and repeat at 4 to 7 day intervals.	Do not apply within 7 days of harvest. Do not apply more than 22.5 lbs (17 lbs active) per acre per crop east of the Mississippi and AR and LA. Do not apply more than 7.5 lbs (5.6 lbs active) per acre per crop west of the Mississippi except AR and LA. Do not feed treated forage to livestock.
(Field and Field Corn for Hybrid Seed Production)				Do not apply within 40 days of harvest. Do not apply more than 15 lbs (11.25 lbs active) per acre per crop. Do not feed treated forage to livestock.
Cranberry	Fruit Rot	3-6	Start applications at mid-bloom and repeat at 7 to 10 day intervals.	Do not apply within 30 days of harvest. Do not apply more than 18 lbs (13.5 lbs active) per acre per season.
Cucumber	Anthraxnose, Cercospora Leaf Spot, Downy Mildew, Gummy Stem Blight, Scab	2-3	Start applications when plants are in the two-leaf stage and repeat at 7 to 10 day intervals. Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces.	Do not apply within 5 days of harvest. Do not apply more than 24 lbs (18 lbs active) per acre per crop.
Fennel	Early Blight, Late Blight	2	Begin in plant beds at emergence. Repeat at 7 to 10 day intervals.	Do not apply within 14 days of harvest. Do not apply more than 16 lbs (12 lbs active) per acre per crop. Do not graze livestock in treated areas.

<b>CROP</b>	<b>DISEASES CONTROLLED</b>	<b>RATE OF MANZATE® PRO-STICK™ PER APPLICATION LBS/ACRE</b>	<b>TIMING/INTERVALS (Also refer to Directions for Use)</b>	<b>RESTRICTIONS/COMMENTS</b>
Grape (East of the Rocky Mountains)	Black Rot, Bunch Rot, Deadarm, Downy Mildew	1.5-4	Apply in sufficient water to provide thorough coverage starting when new shoots are 1/2 to 1 1/2 inches long. Repeat when shoots are 3 to 5 inches long, when shoots are 8 to 10 inches long, and then at 7 to 10 day intervals until fruit is set. For late season control of black rot, deadarm and downy mildew, the use of other approved and recommended fungicides is suggested.	Do not apply within 66 days of harvest. Do not apply more than 24 lbs (18 lbs active) per acre per season.
Grape (West of the Rocky Mountains)	Black Rot, Bunch Rot, Deadarm, Downy Mildew	1.5-2.5	Apply in sufficient water to provide thorough coverage starting when new shoots are 1/2 to 1 1/2 inches long. Repeat when shoots are 3 to 5 inches long, when shoots are 8 to 10 inches long, and then at 7 to 10 day intervals until fruit is set. For late season control of black rot, deadarm and downy mildew, the use of other approved and recommended fungicides is suggested.	Do not apply within 66 days of harvest except in CA where no application can be made after bloom. Do not apply more than 7.5 lbs (5.6 lbs active) per acre per season.
Melon  Cantaloupe, Casaba, Crenshaw, Honeydew, Watermelon	Alternaria Leaf Spot, Anthracnose, Downy Mildew, Gummy Stem Blight, Cercospora Leaf Spot	2-3	Start applications when plants are in the two-leaf stage and repeat at 7 to 10 day intervals. Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces. Some varieties are sensitive to MANZATE® PRO-STICK™ fungicide. Consult State Cooperative Extension Service Specialist prior to use.	Do not apply within 5 days of harvest. Do not apply more than 24 lbs (18 lbs active) per acre per season.
Oat	(See Barley)			
Onion (Dry Bulb), Garlic, Shallot (Furrow Drench)	Botrytis Leaf Blight, Downy Mildew, Neck Rot, Purple Blotch	3	Follow a protective spray schedule starting when diseases are first reported in the area and repeat at 7 day intervals throughout the season.	Do not apply within 7 days of harvest. Do not apply to exposed bulbs. Do not apply more than 30 lbs (22.5 lbs active) per acre per crop.
	Smut	3	Apply 3 lbs per acre as a furrow drench at time of planting onion seeds. Use 75 to 125 gallons of water per acre.	Do not use more than 2.25 lbs active per acre (29,000 linear feet of furrow) with an 18 inch row spacing.
Papaya	Anthracnose (Colletotricum), Phytophthora Fruit Rot, Black Spot (Cercospora)	2.0-2.5 (minimum 50 gals per acre)	Begin at flowering; treat central column crown, blossom area and developing fruit. Repeat at 14 to 21 day intervals.	Do not use more than 35 lbs (26.25 lbs active) per acre per crop. Minimum pre-harvest interval 0 days.
Peanut	Ascochyta Web Blotch, Cercospora Leaf Spot, Rust	1-2	Start application when disease first appears or is reported in area. Repeat sprays at 7 to 14 day intervals. Reduce sprays to a 7 day interval during humid weather.	Do not apply within 14 days of harvest. Do not use more than 16 lbs (12 lbs active) per acre per crop. Do not feed treated vines to livestock.
Pear	(See Pomefruit)			

CROP	DISEASES CONTROLLED	RATE OF MANZATE® PRO-STICK™ PER APPLICATION LBS/ACRE	TIMING/INTERVALS (Also refer to Directions for Use)	RESTRICTIONS/COMMENTS
Pome Fruit  Apple, Pear, Crabapple, Quince	Rusts, Scab, Fabrea Leaf Spot	6.0 Maximum per acre use rate based on thorough coverage dilute sprays. Use 50 gal minimum per acre. Consult State Extension Service if necessary to adjust for variable tree size.	<b>Pre-Bloom/Bloom Use:</b> Begin application at 1/4 to 1/2 inch green tip and continue on a 7 to 10 day schedule through bloom. Use either the “Pre-Bloom/Bloom” or “Extended Application” schedule. <b>DO NOT COMBINE OR INTEGRATE THE TWO TREATMENT SCHEDULES</b>	Do not apply more than 6 lbs (4.5 lbs active) per acre per application. Do not apply after bloom. Do not apply more than 24 lbs (18 lbs active) per acre per year. Do not graze livestock in treated areas. It is recommended that this product be used in an Integrated Pest Management Program.
		3.0 Maximum per acre use rate based on thorough coverage dilute sprays. Use 50 gal minimum per acre. Consult State Extension Service if necessary to adjust for variable tree size.	<b>Extended Application Schedule or for Use in Tank Mixtures:</b> For implementation of IPM programs, applications based on tree-row volume, or for use as a resistance management tool: begin applications at 1/4 to 1/2 inch green tip and continue applications on a 7 to 10 day schedule through the second cover spray. Use either the “Pre-Bloom/Bloom” or “Extended Application” schedule. <b>DO NOT COMBINE OR INTEGRATE THE TWO TREATMENT SCHEDULES</b>	Do not apply more than 3 lbs (2.25 lbs active) per acre per application. Do not apply within 77 days of harvest. Do not apply more than 21 lbs (15.75 lbs active) per acre per year. Do not graze livestock in treated areas. It is recommended that this product be used in an Integrated Pest Management Program.
Potato	Early Blight, Late Blight Black Dot  <b>Disease Suppression:</b> Botrytis	1-2	Begin applications when plants are 4 to 6 inches high by applying 1 lb per acre. As the vines increase in size, apply 1.5 to 2 lbs per acre at intervals of 5 to 10 days or 1 lb per acre at 3 to 5 day intervals.	Do not apply more than 15 lbs (11.2 lbs active) per acre per crop. Do not use within 3 days of harvest in CT, DE, FL, MA, ME, MI, NH, NY, OH, PA, RI, VT, WI, and within 14 days elsewhere. Vine-kill should occur 14 days before harvest. It is recommended that this product be used in an Integrated Pest Management Program.
Potato (Seedpiece Treatment)	Fusarium Decay, Seedborne Common Scab	1.25 per 50 gal	Dip whole or cut potato tubers in 1.25 lbs MANZATE® PRO-STICK™ fungicide per 50 gallons of water. Place treated tubers in a clean container following treatment and plant as soon as possible. Spread treated seedpieces in a cool place if held before planting.	Do not use treated seed potatoes for food or feed purposes.
Squash (Summer Squash, Including Edible Gourd)	Downy Mildew	2-3	Start applications when plants are in the two-leaf stage and repeat at 7 to 10 day intervals. Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces.	Do not apply within 5 days of harvest. Do not apply more than 24 lbs (18 lbs active) per acre per crop.
Sugar Beet	Cercospora Leaf Spot	1.5-2	Begin when disease first threatens. Repeat at 7 to 10 day intervals.	Do not apply within 14 days of harvest. Do not apply more than 14 lbs (10.5 lbs active) per acre per crop. Do not feed treated sugar beet tops to livestock.

<b>CROP</b>	<b>DISEASES CONTROLLED</b>	<b>RATE OF MANZATE® PRO-STICK™ PER APPLICATION LBS/ACRE</b>	<b>TIMING/INTERVALS (Also refer to Directions for Use)</b>	<b>RESTRICTIONS/COMMENTS</b>
Tomato (East of the Mississippi River)	Anthracnose, Early Blight, Gray Leaf Spot, Late Blight, Leaf Mold, Septoria Leaf Spot	0.75-1.5	Start application when seedlings emerge or transplants are set. Repeat at 3 to 7 day intervals throughout the season.	Do not apply within 5 days of harvest. Do not apply more than 22.4 lbs (16.8 lbs active) per acre per crop.
	Bacterial Speck and Spot	1.5-3	Start application when seedlings emerge or transplants are set. Repeat at 7 to 10 day intervals throughout the season.	Do not apply within 5 days of harvest. Do not apply more than 22.4 lbs (16.8 lbs active) per acre per crop. Use a full rate of a fixed copper fungicide in tank mix combination with a half to full rate of MANZATE® PRO-STICK™. Follow the application interval recommended on the copper fungicide label.
Tomato (West of the Mississippi River)	Anthracnose, Early Blight, Gray Leaf Spot, Late Blight, Leaf Mold, Septoria Leaf Spot	0.75-1.0	Start application when seedlings emerge or transplants are set. Repeat at 3 to 7 day intervals throughout the season.	Do not apply within 5 days of harvest. Do not apply more than 8.5 lbs (6.4 lbs active) per acre per crop.
	Bacterial Speck and Spot	1.5-2	Start application when seedlings emerge or transplants are set. Repeat at 7 to 10 day intervals throughout the season.	Do not apply within 5 days of harvest. Do not apply more than 8.5 lbs (6.4 lbs active) per acre per crop. Use a full rate of a fixed copper fungicide in tank mix combination with a half to full rate of MANZATE® PRO-STICK™. Follow the application interval recommended on the copper fungicide label.
Watermelon	(See Melon)			
Wheat (Including Triticale)	(See Barley)			

## SEED TREATMENTS

Users making commercial seed applications must follow provisions within the Non-Agricultural Use Requirements Box. Users conducting seed treatments on agricultural establishments must follow provisions within the Agricultural Use Requirements Box.

Seed to be treated must be clean and well-cured prior to treatment. Prior to seed treatment, a dye must be added to the treating slurry so that an unnatural color will distinguish the seed as treated.

MANZATE® PRO-STICK™ may be applied to dry seed with conventional slurry or mist seed treating equipment, or as a plant-box application. For best results, seed must be covered uniformly with fungicide.

LABEL TREATED SEED: “Do not use for food, feed or oil purposes. This seed treated with MANZATE® PRO-STICK™ fungicide.”

CROP	DISEASES	SEED TREATMENT RATE- APPLY AS A SLURRY OZ./BU.	OZ./100 LBS.
Barley	Bunt, Covered Smut, Damping-Off, Fake Loose Smut, Seed Decay, Seedling Blights	1.3 to 2.0	2.7 to 4.2
Corn	Damping-Off, Seed Rot, Seedling Blights	1.5 to 3.0	2.7 to 5.4
Cotton Acid Delinted	Damping-Off, Seedling Blights	--	3.0
Cotton Reginned	Damping-Off, Seedling Blights	--	6.0
Flax	Seed Decay, Seedling Blights, Damping-Off	2.0 to 4.0	3.6 to 7.1
Oat	Damping-Off, Seedling Blights, Seed Decay, Smuts	1.3 to 2.0	4.0 to 6.3
Peanut (Shelled)	Damping-Off, Seed Rots, Seedling Blights	2.0 to 4.0	8.0 to 16.0
Rice	Achyla, Other Soil and Seedborne Fungi Causing Seed Rot and Reduced Seedling Vigor	--	2.0 to 4.0 Apply before, during or after soaking in water
Rye	Bunt, Covered Smut, Damping-Off, Seed Decay, Seedling Blights	1.3 to 2.0	2.3 to 3.6
Safflower	<i>Puccinia carthami</i> (Which Causes Foot-and-Rot Disease and Foliage Rust Disease)	--	2.0
Sorghum	Covered Kernel Smut, Damping-Off, Seedling Blights, Seed Rots	1.5 to 2.5	2.7 to 4.5
Tomato	Damping-Off, Seedling Blights, Seed Rots	--	8.0
Wheat (including Triticale)	Bunt, Covered Smut, Damping-Off, Seed Decay, Seedling Blights	1.3 to 2.0	2.2 to 3.3



## STORAGE AND DISPOSAL

**PESTICIDE STORAGE: Important-**Never allow MANZATE® PRO-STICK™ to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of MANZATE® PRO-STICK™ as a fungicide and create vapors which may be flammable. Keep container closed when not in use. Store product in original container only, away from other pesticides, fertilizer, food or feed.

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**ATTENTION:** This product contains mancozeb and ETU, chemicals known to the State of California to cause cancer in laboratory animals. ETU is also known to the State of California to cause birth defects or other reproductive harm in laboratory animals.

## IMPORTANT INFORMATION READ BEFORE USING PRODUCT

### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorus, Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorus, Inc. and Seller harmless for any claims relating to such factors.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UNITED PHOSPHORUS, INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.**

To the extent consistent with applicable law, United Phosphorus, Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHORUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED PHOSPHORUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorus, Inc.

Manzate is a registered trademark of United Phosphorus, Inc.

© 2010 United Phosphorus, Inc. All rights reserved.

A01397571

70506-234(082710-3812)