

ONSET[®] 3.6L

Fungicide for the control of specified diseases on asparagus, barley, beans, corn, cotton, cucurbit vegetables, grasses grown for seed, hops, dry bulb onion and garlic, green onions, leafy Brassica greens, garden beet roots and tops, lychee, okra, pecan, peanut, soybean, sunflower, turnip, and wheat.

ACTIVE INGREDIENT:

Tebuconazole,

alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol 38.7%

OTHER INGREDIENTS: 61.3%

TOTAL: 100.0%

Contains 3.6 pounds tebuconazole per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	
Note To Physician: No specific antidote. Treat symptomatically. Symptoms of Poisoning: The compound does not cause any definite symptoms that would be diagnostic.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist. Contact with the eyes may cause irritation.

EPA Reg. No. 1381-203

EPA Est. No. 42750-MO-01

Manufactured For:

Winfield Solutions, LLC

P.O. Box 64589

St. Paul, MN 55164-0589

NET CONTENTS: 2.5 Gals. (9.46 Liters)

LOT. NO. _____

1/1003/2

WINFIELD[™]

AgriSOLUTIONS[™]

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

1. Coveralls
2. Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
3. Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Use label language appropriate for container size and type.

Nonrefillable containers. Do not reuse or refill this container. Clean container promptly after emptying.

Nonrefillable container equal to or less than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable container greater than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident, call
CHEMTREC 1-800-424-9300.**

SPRAY VOLUME

ONSET 3.6L may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

CHEMIGATION

Apply ONSET 3.6L through irrigation equipment only to crops and diseases for which the chemigation use is specified. Apply this product only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

Mixing: Add specified amount of ONSET 3.6L into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the ONSET 3.6L should be thoroughly dispersed prior to the addition of other materials.

Compatibility: To determine the compatibility of ONSET 3.6L with other products, the following procedure should be followed: Pour the specified proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF ONSET 3.6L
Asparagus	Rust (Puccinia spp.)	4 to 6 fl. oz per acre
<p>Notes: Apply ONSET 3.6L as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl oz of ONSET 3.6L per acre (0.11 lb ai - 0.17 lb ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not apply within 100 days harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl oz/acre or 0.51lb ai/acre).</p> <p>General Comments: Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with ONSET 3.6L. ONSET 3.6L is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating ONSET 3.6L with other DMI fungicides may lead to resistance.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF ONSET 3.6L
Barley	Rusts (Puccinia spp.) Head blight (Fusarium spp.) Suppression	4 fl oz per acre
<p>Notes and Restrictions:</p> <ul style="list-style-type: none"> • Apply ONSET 3.6L in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. • A maximum of 4 fl oz of ONSET 3.6L may be applied per acre per crop season. • Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. • Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of ONSET 3.6L. • Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. • Restricted-entry interval (REI) = 12 hours. 		
<p>Application Timing Directions:</p> <ul style="list-style-type: none"> • Rusts: Apply at the earliest sign of rust pustules on foliage. • Fusarium head blight: Optimal timing for head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants. 		
<p>General Comments:</p> <ul style="list-style-type: none"> • For optimum disease control, the lowest specified rate of a spray surfactant should be tank mixed with ONSET 3.6L. • ONSET 3.6L must have 2 to 4 hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. • ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3). 		

CROP	DISEASE	RATE OF ONSET 3.6L
Beans (fresh & dry except succulent shelled)	Rust (Uromyces appendiculatus)	4 to 6 fl. oz per acre
Notes: Apply ONSET 3.6L in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. Beans, fresh: ONSET 3.6L may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of ONSET 3.6L per acre per crop season. Beans, dry: ONSET 3.6L may be applied up to 14 days before harvest. Do not apply more than 12 fl. oz. of ONSET 3.6L per acre per crop season.		
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with ONSET 3.6L. ONSET 3.6L must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3).		
Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF ONSET 3.6L
Corn (sweet corn, field corn, field corn grown for seed, and popcorn)	Rust (Puccinia spp.) Northern leaf blight (Helminthosporium turcicum) Southern leaf blight (Helminthosporium maydis) Northern leaf spot (Helminthosporium carbonum) Gray leaf spot (Cercospora zeae-maydis)	4 to 6 fl. oz per acre
Notes: Apply ONSET 3.6L in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. A maximum of 24 fl. oz. (1.5 pint) of ONSET 3.6L may be applied per acre per crop season. Sweet corn: ONSET 3.6L may be applied up to 7 days before harvest of ears or forage, and 49 days before the harvest of fodder. Field, seed or popcorn: ONSET 3.6L may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.		
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with ONSET 3.6L. ONSET 3.6L must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3).		
Restricted-entry interval (REI) for sweet corn = 19 days.		
Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.		

CROP	DISEASE	RATE OF ONSET 3.6L
Cotton	Southwestern cotton rust (Puccinia cacabata)	6 to 8 fl. oz per acre
Notes: Apply ONSET 3.6L in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. ONSET 3.6L may be applied up to 30 days before harvest. Do not apply more than 24 fl. oz. of ONSET 3.6L per acre per crop season.		
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with ONSET 3.6L. ONSET 3.6L must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3).		
Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF ONSET 3.6L
Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd, (includes hyotan, cucuzza, hechima and Chinese okra) Momordica spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternet squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon	Powdery mildew (Sphaerotheca fuliginea / Podosphaera xanthii) (Erysiphe cichoracearum)	4 to 6 fl. oz per acre
	Gummy stem blight – suppression (Didymella bryonae) (watermelon, squash, pumpkin, and melons only)	8 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. ONSET 3.6L may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of ONSET 3.6L per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with ONSET 3.6L. ONSET 3.6L must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3).		
Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF ONSET 3.6L
Grasses Grown for Seed	Rusts (Puccinia spp.)	4 to 8 fl oz per acre
	Apply the specified rate of ONSET 3.6L as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	
	Powdery mildew	4 to 8 fl oz per acre
	Apply specified rate of ONSET 3.6L when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	

General Comments: Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.

For optimum benefit, the lowest labeled rate of a spray surfactant should be tank mixed with ONSET 3.6L.

A maximum of 16 fluid ounces (1 pint) may be applied per acre per crop season. ONSET 3.6L may be applied up to 4 days before harvest. Chaff, screenings and straw from treated areas may be used for feed purposes; however, do not forage, cut green crop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.

Restricted-entry interval (REI) = 12 hours.

CROP	DISEASE	RATE OF ONSET 3.6L
Hops	Powdery mildew (<i>Sphaerotheca humuli</i> / <i>Sphaerotheca macularis</i>)	4 to 8 fl. oz per acre
Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. ONSET 3.6L may be applied up to 14 days before harvest. Do not apply more than 32 fl. oz. of ONSET 3.6L per acre per crop season. Increase the spray volume and the application rate as vine growth increases during the season.		
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with ONSET 3.6L. ONSET 3.6L must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3).		
Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF ONSET 3.6L
Dry bulb onion Garlic Great-headed (elephant) garlic Welch onion Shallot	White rot (<i>Sclerotium cepivorum</i>)	White rot: 20.5 fl oz per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot.
	Rust (<i>Puccinia allii</i> , <i>Puccinia porri</i>)	4 to 6 fl. oz. per acre
	Purple blotch (<i>Alternaria porii</i>)	
<p>White rot: For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz. ONSET 3.6L per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl oz/acre.</p> <p>Rust: For the control of rust make foliar applications at the rate of 4 to 6 fl. oz ONSET 3.6L per acre per application. Repeat at an interval of 10 to 14 days. Apply ONSET 3.6L in a protective spray schedule or when weather conditions are favorable for rust development.</p> <p>Notes: Do not apply more than 32.5 fl. oz. of ONSET 3.6L per acre per crop season if an in-furrow treatment is made. If ONSET 3.6L is not applied as an in-furrow treatment then do not apply more than 12 fl oz. ONSET 3.6L per acre per season as a foliar spray. Do not apply within 7 days of harvest (PHI = 7 days).</p> <p>General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with ONSET 3.6L. ONSET 3.6L must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF ONSET 3.6L
Green Onion Leek Spring Onion Scallion Japanese bunching onion Green shallots Green eschalots	White rot (Sclerotium cepivorum) suppression only Rust (Puccinia allii, Puccinia porri) Purple blotch (Alternaira porii)	4 to 6 fl. oz per acre
For the control of diseases make foliar applications using an interval of 10 to 14 days. Apply ONSET 3.6L in a protective spray schedule or when weather conditons are favorable for rust development. Notes: Do not apply more than 24 fl. oz. of ONSET 3.6L per acre per crop season. Do not apply within 7 days of harvest (PHI = 7 days).		
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with ONSET 3.6L. ONSET 3.6L must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entrv interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF ONSET 3.6L
Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuma Mustard greens Mustard spinach Rape greens Turnip greens	Cercospora leaf spot (<i>Cercospora brassicicola</i>) Powdery mildew (<i>Erysiphe cruciferarum</i>) Alternaria leaf spot (<i>Alternaria brassicicola</i>)	3 to 4 fl. oz per acre
<p>Notes: Do not apply more often than once every 10 days. Do not apply more than 16 fl. oz. of ONSET 3.6L per acre per crop season. Do not apply within 7 days of harvest (PHI = 7 days)</p> <p>General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant should be tank-mixed with ONSET 3.6L. ONSET 3.6L must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3). Restriction: Application to turnip greens is limited to East of the Rockies.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF ONSET 3.6L
Garden beet roots and tops (leaves)	Cercospora leaf spot (<i>Cercospora beticola</i>)	3 to 7.2 fl. oz per acre
<p>Notes: Make applications on a 14 day intervals. Do not apply more than 28.8 fl. oz. ONSET 3.6L per acre season. Do not apply within 7 days of harvest (PHI = 7 days).</p> <p>General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant should be tank-mixed with ONSET 3.6L. ONSET 3.6L must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF ONSET 3.6L
Lychee	Anthrachnose (<i>Colletotrichum gloesporioides</i>)	4 to 6 fl. oz per acre
<p>Notes: Begin first application of ONSET 3.6L as panicle emerges. Spray up to 6 fl. oz. per acre every 10 days thereafter for a total of 8 sprays. Apply specific dosage in a minimum of 50 gallons of spray solution per acre by ground only. Do not apply more than 48 fl. oz. of ONSET 3.6L per acre season. ONSET 3.6L can be applied up to and including the day of harvest (PHI = 0 days).</p> <p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with ONSET 3.6L. ONSET 3.6L must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 2 days.</p>		

CROP	DISEASE	RATE OF ONSET 3.6L
Okra	Cercospora leaf spot (Cercospora spp.)	4 to 6 fl. oz per acre
<p>Notes: Apply specified dosage of ONSET 3.6L in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Applications may be made no closer than 3 days before harvest. Do not apply more than 24 fl. oz. of ONSET 3.6L per acre per season.</p> <p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with ONSET 3.6L. ONSET 3.6L must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF ONSET 3.6L
Peanut	SOILBORNE: Sclerotium stem and pod rot (white mold, southern blight, southern stem rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only) FOLIAR: Early leaf spot Late leaf spot Leaf rust Web blotch (Phoma) Pepper spot (Leptosphaerulina)	7.2 fl oz per acre
<p>FOUR-APPLICATION SPRAY PROGRAM: Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Applications of chlorothalonil should be made prior to and following applications of ONSET 3.6L to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label recommended rate of a spray surfactant should be tank-mixed with ONSET 3.6L.</p> <p>LEAF SPOT ADVISORY SCHEDULE: For control of soilborne diseases in an advisory schedule, apply ONSET 3.6L in the first advisory spray in July and continue ONSET 3.6L applications at 14-day intervals. Applications after August 15 should be tank mixed with chlorothalonil for resistance management purposes.</p>		

GENERAL DIRECTIONS: For optimum control of the specified soilborne diseases, four consecutive applications of ONSET 3.6L must be made at 14-day intervals.

A maximum of 28.8 fluid ounces of ONSET 3.6L may be applied per crop season. ONSET 3.6L may be applied up to 14 days before harvest. Do not feed hay or threshings or allow livestock to graze in treated areas.

ONSET 3.6L is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with ONSET 3.6L as a leaf spot resistance management strategy. A spray surfactant is not necessary when ONSET 3.6L is tank mixed with chlorothalonil. Mixing or alternating ONSET 3.6L with other DMI fungicides may lead to resistance.

ONSET 3.6L must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by *Sclerotium rolfsii* and *Rhizoctonia solani*. Drought conditions will decrease the effectiveness of ONSET 3.6L against the root and pod rots.

Use ONSET 3.6L in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices.

Timing of ONSET 3.6L Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot

Spray Program	ONSET 3.6L Application No.	Chlorothalonil Application No.
7 applications	3, 4, 5 and 6	1, 2 and 7

Restricted-entry interval = 12 hours.

CROP	DISEASE	RATE OF ONSET 3.6L
Pecan	Brown leaf spot (<i>Sirosporium diffusum</i>) Downy spot (<i>Mycosphaerella caryigena</i>) Liver spot (<i>Gnomonia caryae</i>) Scab (<i>Cladosporium caryigenum</i>) Vein spot (<i>Gnomonia nerviseda</i>) Zonate leaf spot (<i>Grovesinia pyramidalis</i>)	4 to 8 fl. oz per acre

Notes: Apply ONSET 3.6L in a preventive spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. ONSET 3.6L may be applied at 4 fl. oz. per acre in tank-mix with the labeled rate of Super-Tin® in cover sprays. Follow label directions for the use of Super-Tin. Do not add a surfactant to the spray solution when tank-mixing ONSET 3.6L with Super-Tin. Apply ONSET 3.6L in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. per acre of ONSET 3.6L to full-size mature trees, and 4 to 6 fl. oz. per acre of ONSET 3.6L to smaller trees. Apply the high rate to

varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases. Do not apply after shucks begin to split. A maximum of 32 fl. oz. of ONSET 3.6L may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with ONSET 3.6L. ONSET 3.6L must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.

Restricted-entry interval (REI) = 12 hours.

CROP	DISEASE	RATE OF ONSET 3.6L
Soybean	Rust (<i>Phakopsora pachyrhizi</i>) Powdery mildew (<i>Microsphaera diffusa</i>)	3 to 4 fl. oz per acre
<p>Use Directions: Apply ONSET 3.6L as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. The lowest labeled rate of a spray surfactant must be tank-mixed with ONSET 3.6L. ONSET 3.6L should be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment. Restrictions: Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply more than 12 fl. oz./A per use season.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF ONSET 3.6L
Sunflower	Rust (<i>Puccinia helianthi</i>)	4 to 6 fl. oz per acre
<p>Notes: Apply specified dosage of ONSET 3.6L at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Apply specific dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 16 fl. oz. of ONSET 3.6L per acre per crop season or within 50 days of harvest.</p> <p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with ONSET 3.6L. Contact your state Extension Service or Winfield Solutions representative for a list of approved surfactants. ONSET 3.6L must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF ONSET 3.6L
Turnip (Application is limited to East of the Rockies)	<i>Cercospora</i> leaf spot (<i>Cercospora brassicicola</i>)	4 to 7.2 fl. oz per acre
<p>Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. ONSET 3.6L may be applied up to 7 days before harvest. Do not apply more than</p>		

28.8 fl. oz. of ONSET 3.6L per acre per crop season.

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with ONSET 3.6L. ONSET 3.6L must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering. ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3).

Restriction: Application to turnip greens is limited to East of the Rockies.

Restricted-entry interval (REI) = 12 hours.

CROP	DISEASE	RATE OF ONSET 3.6L
Wheat	Rust leaf, stem and stripe (Puccinia spp.) Head blight or scab (Fusarium spp.) - Suppression	4 fl oz. per acre
<p>Notes and Restrictions:</p> <ul style="list-style-type: none">Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.Apply ONSET 3.6L in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.A maximum of 4 fl oz of ONSET 3.6L may be applied per acre per crop season.Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding.Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with ONSET 3.6L.Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.Restricted-entry interval (REI) = 12 hours.		
<p>Application Timing Directions:</p> <ul style="list-style-type: none">Rusts: Apply at the earliest sign of rust pustules on foliage.Fusarium head blight: Optimal timing for head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).		
<p>General Comments:</p> <ul style="list-style-type: none">For optimum disease control, the lowest specified rate of a spray surfactant should be tank mixed with ONSET 3.6L.ONSET 3.6L must have 2 to 4 hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering.ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3).		

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

Apply only during alternate years in fields adjacent to aquatic areas listed above.

Do not apply by ground or air within 100 feet of aquatic areas listed above.

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Spray Drift Management: For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

AgriSolutions™ is a trademark and ONSET® is a registered trademark of Winfield Solutions, LLC.