CROP TOLERANCE: Quali-Pro Oxadiazon 50 WSB may be used as an over-the-top spray only on the ornamental species listed

VARIETIES SPECIES Ice Plant Large Leaf English, Algerian Daisv African Trailing Periwinkle Major Minor

TANK MIX COMBINATIONS: Ornamental Plantings, Noncropland

Tank mix combinations consisting of Quali-Pro Oxadiazon 50 WSB with Roundup® are recommended to control existing undesirable vegetation. When applied as a directed spray, Quali-Pro Oxadiazon 50 WSB plus Roundup will provide pre- and postemer- plants. gence control of the susceptible species that are listed on the respective labels of the two herbicides.

> BROADCAST APPLICATION RATES: OXADIAZON/ROUNDUP TANK MIX

Existing Broadleaf and Annual Weeds	Quali-Pro Oxadiazon 50 WSB Number of Water-Soluble Packets per Acre	Roundup Quarts Per Acre
Less Than 6 Inches Tall	4 – 8	1
Greater Than 6 Inches Tall	4 – 8	1 1/2
Existing Perennials	4 – 8	2 – 5

These rates are intended as general information only. Consult the rate sections in individual product labels for detailed information regarding specific weed species controlled and rates of application.

Review the instructions under Application Rates in the GENER-AL INFORMATION section above for additional application infor-

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not contaminate water, food or feed by storage or disposal. Do not store in or around the home. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Nonrefillable Container (flexible-bag-all weights):

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, Dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (rigid-fifty lbs. or less): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two

MIXING INSTRUCTIONS: Quali-Pro Oxadiazon 50 WSB should be added to the tank and agitated before adding Roundup. Please refer to the Roundup label for information on the use of surfactants. Only mix the amount of spray needed for immediate application needs, as settling can occur if the spray mixture sits over an extended period of

PRECAUTION: Roundup is not recommended as an over-the-top spray on ornamentals. Exercise extreme care to avoid contact of spray containing Roundup with foliage, green bark, stems, or bare roots of desirable turfgrasses, shrubs, trees, or other desirable vegetation since severe damage or destruction may result. If spraying a tank mix with Roundup on areas adjacent to desirable plants, use a shield made of plyboard, cardboard, or sheet metal while spraying to help prevent spray from contacting foliage of desirable

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following CONDITIONS. DISCLAIMER OF WARRANTIES. and LIMITATIONS OF LIABILITY

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer. DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America. Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special. incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replace-

Quali-Pro is a trademark of Makhteshim Agan of North America.

ment of product.

Roundup is a registered trademark of Monsanto Agricultural Company



Oxadiazon 50 WSB Herbicide

A pre-emergent herbicide for the control of broadleaf weeds and annual grasses in turf and ornamentals

FOR SALE TO AND USE BY PROFESSIONAL APPLICATORS ONLY, NOT FOR SALE TO OR USE BY HOMEOWNERS/CONSUMERS. NOT FOR USE ON HOME LAWNS.

ACTIVE INGREDIENT	% BY WT.	
Oxadiazon: 3-[2,4-dichloro-5-(1-methylethoxy)phenyl]-	
5-(1,1-dimethylethyl)-1,3,4-oxadiazol-2(3H)-one	50.0%	
NERT INGREDIENTS:	<u>50.0%</u>	
TOTAL:	100.0%	

EPA Reg. No. 66222-209

14158

EPA Est. No. 65387-AR-001

KEEP OUT OF REACH OF CHILDREN **WARNING / AVISO**

Si usted no entiende la etiqueta, busque a alquien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

For additional precautionary, handling, and use statements, see inside of this booklet.

Manufactured for:

Makhteshim Agan of North America, Inc. 4515 Falls of Neuse Road, Suite 300 Raleigh, NC 27609

Note to Physician: There is no specific antidote. Appropriate supportive and symptomatic treatment as indicated by the patient's condition is recommended.

IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Riinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor treatment advice.
IF IN EYES:	Hold eye open and rinse slowly and ge with water for 15 – 20 minutes. Remove contact lenses, if present, afte the first 5 minutes; then continue rinsin Call a poison control center or doctor to treatment advice.
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or a ambulance; then give artificial respiration preferably by mouth-to-mouth if possible Call a poison control center or doctor for further treatment advice.
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if abl swallow. Do not induce vomiting unless told to be poison control center or doctor.

FIRST AID

EPA 011008/Trans 010410/Notif 030510/Rev D

• Do not give anything by mouth to an

unconscious person.

Have the product container or label with you when calling a poison

control center or doctor or going for treatment. You may also con-

tact Prosar at 1-877-250-9291 for emergency medical treatment

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes substantial but temporary eye injury. Causes skin irritation. Harmful if inhaled or absorbed through skin. Do not get in eyes, on skin, or on clothing. Avoid breathing dust,

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are barrier laminate. neoprene rubber (>14 mils), and polyethylene. If you want more options, follow the instructions for Category A on an EPA chemical-resistant category selection chart.

Mixers and loaders must wear:

- · Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of waterproof materials
- Chemical-resistant apron

All other handlers must wear:

- · Coveralls over short-sleeved shirt and short pants
- · Socks and chemical-resistant footwear · Chemical-resistant gloves made of waterproof materials
- Chemical-resistant headgear for overhead exposure

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

See ENGINEERING CONTROLS STATEMENTS for additional require-

ENGINEERING CONTROLS STATEMENTS

Water-soluble packets, when used correctly, qualify as a closed mixing/loading system under the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240 (d) (4)]. Mixers and loaders using water-soluble packets must:

- · Wear the personal protective equipment above for mixers/loaders,
- Be provided and have immediately available for use in an emergency, such as a broken package, spill, or equipment breakdown: chemical-resistant footwear and a NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any N. R. P. or HE filter.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

Quali-Pro® Oxadiazon 50 WSB should not be used in areas where irrigation or rainfall results in the direct or indirect (via ditches and canals) contamination of surface waters through dissolved runoff or erosion of soil particles to which oxadiazon is absorbed. Do not apply this product within 15 feet of bodies of water that may serve as sources of drinking water.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner incon-

Read entire label before using this product. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water.

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material

NON-AGRICULTURAL USE REQUIREMENTS

The requirement in this section applies to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

GENERAL INFORMATION Not for use on home lawns.

One water-soluble packet is equivalent to 1 pound of product, which equals 1/2 pound of active ingredient.

Quali-Pro Oxadiazon 50 WSB is a preemergent herbicide for control of many broadleaf weeds and annual grasses in turf and on a variety of woody ornamental shrubs and trees. Quali-Pro Oxadiazon 50 WSB controls weeds by killing the seedlings of young weeds as they come in contact with the herbicide during germination.

> Make applications with a properly calibrated sprayer that will apply The state of Arizona has not approved this product for use on agrithe spray solution uniformly. Uniform application is essential for cultural sites. Do not use this product on uses considered by the satisfactory weed control. Calibrate the sprayer before using, Arizona statutes to be agricultural uses. according to the manufacturer's directions, and check the equip-

> Application Rates: Do not use more than 12 pounds (12 packets) Augustinegrass, and zovsia turf such as fairways, golf courses. of product per acre per year except in areas where there is heavy parks, and non-residential lawns. Quali-Pro Oxadiazon 50 WSB weed infestation. In areas of heavy weed infestation, the maximum controls crabgrass, annual bluegrass, goosegrass, Virginia buttonweed, and annual sedge. Do not use on zovsia turf or St. application rate is 16 pounds (16 packets) of product per acre per year. Do not apply more than 8 pounds (8 packets) of product per Augustinegrass in California. acre per application.

> Wherever a range of application rates are listed on this label (for greenup. Rake turf area to remove leaves and trash. If necesexample: "4-6 water-soluble packets"), use higher rates of Qualisarv, mow turf to ensure the spray solution reaches the soil sur-Pro Oxadiazon 50 WSB where heavy weed pressure is anticipatface. Measure the area to be treated and apply the amount ed, or where medium- or fine-textured soils exist. Soil textures are required uniformly. If rain is not expected shortly after application defined as follows: of this product, sprinkle the turf thoroughly.

- Coarse: Sand, sandy loam, loamy sand
- . Medium: Loam, silt loam, silt

ment daily to assure that it is working properly.

 Fine: Sandy clay loam, silty clay loam, clay loam, sandy sod. Ensure that the spray reaches the soil surface. If rain is not clay, silty clay, clay

Lower rates may be used on light (coarse) texture, low organic matter (< 1.0% OM) soils; however, these guidelines are intended as general information only. Consult the list of annual grasses and broadleaf weeds below for rates used to control specific weeds.

When a range of spray volumes is recommended, use a sufficient amount to ensure adequate spray coverage under your use conditions. Generally, larger plants require larger volumes.

GENERAL HANDLING INSTRUCTIONS

Each Quali-Pro Oxadiazon 50 WSB water-soluble bag is sealed within a plastic outer sleeve. Do not break the seal until ready to use. Open the end of the outer sleeve by tearing it along the print ed solid line. Note: Do not use a knife or other sharp object to break the seal. Once the outer protective sleeve is opened, the water-soluble bag inside can be dropped into the spray tank at the appropriate time without touching the bag.

HANDLING PRECAUTIONS

- Do not allow the empty outer sleeve or the tear-strip to fall into the spray tank. They are not water soluble and could clog the sprayer.
- Do not handle the inner bag with wet gloves, as the bag will begin to dissolve on contact. This can occur with even a small amount
- . Do not open the inner bags, and do not attempt to remove the product from them. The water-soluble packet is not to be subdivided or re-measured.
- Do not open the PVA bag. A partial bag should not be used for a

PREEMERGENCE WEED CONTROL

Use the entire contents of the water-soluble bag.

For use in dormant, established bermudagrass, St.

State Specific Restrictions:

thoroughly.

Do not permit domestic animals to graze treated areas. Avoid con-

Apply Quali-Pro Oxadiazon 50 WSB at least 2-3 weeks prior to

Hawaii recommendation only: Quali-Pro Oxadiazon 50 WSB may

be applied to newly sprigged zoysiagrass that is being grown for

expected shortly after application of this product, irrigate the turf

tamination of water that is intended for irrigation and domestic use.

WEEDS CONTROLLED	NUMBER OF QUALI-PRO OXADIAZON 50 WSB WATER-SOLUBLE PACKETS PER ACRE	INSTRUCTIONS
Annual Bluegrass (Poa annua)* Crabgrass (Digitaria spp.) Goosegrass (Eleusine indica) Annual Sedge (Cyperus compressus)*	4 - 6	Apply in late winter or early spring before weed seed germination and turf greenup. NOTE: Treatment is effective only against spring germinating <i>Poa annua</i> .
Virginia Buttonweed (Diodia virginiana)	6	Apply in 30 – 150 gallons of water per acre.

^{*} Not currently registered in California

Review the instructions under Application Rates in the GENERAL INFORMATION section above for additional application information.

ANNUAL WEED CONTROL FOR BERMUDAGRASS ESTABLISHMENT FROM SPRIGS

		Chambaudanuuaa1	
WEEDS CONTROLLED	NUMBER OF QUALI-PRO OXADIAZON 50 WSB WATER-SOLUBLE PACKETS PER ACRE	INSTRUCTIONS	Shepherdspurse ¹ Speedwell Sow Thistle ¹ Spotted Catsear
rabgrass, Annual Sedge, and posegrass	4 - 6	Apply immediately before or immediately after sprigging. Disturbance of the soil surface after applying Quali-Pro Oxadiazon 50 WSB may disrupt the herbicide barrier and reduce product	Swinecress* Yellow Woodsorrel
		effectiveness. For best results, irrigate immediately after application.	COMMON NAME Annual Sedge ¹ Barnvardgrass*

Quali-Pro Oxadiazon 50 WSB must not be applied to newly seeded areas. Wait until 4 months after treatment before seeding into treated areas. Do not use Quali-Pro Oxadiazon 50 WSB on putting greens or tees because of the varying cultural practices used. Apply this product in 30 - 150 gallons of water per acre.

Review the instructions under Application Rates in the GENER-AL INFORMATION section above for additional application infor-

ORNAMENTALS AND CONIFER NURSERIES

TREE, WOODY ORNAMENTAL SHRUB AND CONIFER NURS-ERY APPLICATIONS: Apply Quali-Pro Oxadiazon 50 WSB before weed seed germination as a directed spray to the soil surface or as an over-the-top spray only on species listed in this label. Quali-Pro Oxadiazon 50 WSB can be applied to both established and newly transplanted plants. Apply to actively growing or dormant plants. However, Quali-Pro Oxadiazon 50 WSB should not be applied during bud break or during the 4 weeks immediately following bud

Rainfall or overhead irrigation (one-half inch) should follow immediately after application of Quali-Pro Oxadiazon 50 WSB to improve weed control activity. Rainfall or overhead irrigation imme-

diately after over-the-top sprays is necessary to move the Quali-Pro Oxadiazon 50 WSB from the foliage to the soil surface.

Do not mix Quali-Pro Oxadiazon 50 WSB into the soil. Cultivating or disturbing the soil may reduce the weed control activity of Quali-Pro Oxadiazon 50 WSB. Do not apply in greenhouses as plant injury may occur due to codistillation of the active ingredient onto leaf surfaces in condensation.

USE RATES: Ground application – Apply Quali-Pro Oxadiazon 50 WSB to ornamentals with a broadcast spreader in a minimum of 50 gallons per acre at 4 to 8 water-soluble packets per acre, depending on the weeds to be controlled, as listed below. Remove existing weed growth before application. Review the instructions under Application Rates in the GENERAL INFORMATION section above for additional application information.

Quali-Pro Oxadiazon 50 WSB at 4 to 8 water-soluble packets per acre is effective in controlling broadleaf weeds and annual grasses such as the following:

ANNUAL BROADLEAF WEEDS

Liverwort**

Petty Spurge

Prostrate Spurge**

Redroot Pigweed

Carpetgrass

Fall Panicum

Goosegrass*

Green Foxtail

Wild Oats1

Ripgut Bromegrass¹

Sow Thistle, and Wild Oats.

Crabgrass

Pennsylvania Smartweed

COMMON NAME SCIENTIFIC NAME Arborvitae (Cardamine spp.) Baby's Breath Bittercress Bristly Oxtongue¹ (Picris echiodes) Barberry (Mollugo verticillata) Carpetweed Boxwood Cheeseweed (Malva parviflora) Cedar Common Groundsel* (Senecio vulgaris) Cranberry Common Purslane (Portulaca oleracea) Dodonaea (Oenothera spp.) Evening Primrose* Euonymus Fiddleneck1 (Amsinckia intermedia) ireweed¹ (Epilobium angustifolium) False Cypress Florida Puslev (Richardia scabia) Galinsoga* (Galinsoga spp.) Golden Ragwort (Senecio aureus) Lambsquarters Gladiolus (Chenopodium album)

(Marchantia spp.)

(Euphorbia peplus)

(Euphorbia supine)

(Veronica spp.)

(Oxalis stricta)

ANNUAL GRASSES

*Use 6 to 8 water-soluble packets of this product per acre.

Not currently registered in California for use on Annual

Bluegrass, Annual Sedge, Bristly Oxtongue, Cheeseweed,

Fiddleneck, Fireweed, Ripgut Bromegrass, Shepardspurse,

PLANT TOLERANCE: Quali-Pro Oxadiazon 50 WSB may be used

as a directed spray to the soil surface on a wide variety of field-

grown and container-grown woody ornamental shrubs and trees

and on conifers in nurseries. Quali-Pro Oxadiazon 50 WSB may be

used as an over-the-top spray in a minimum of 50 gallons per acre

at the recommended rates only on the ornamental species below:

**Use 8 water-soluble packets of this product per acre.

(Sonchus oleraceus)

(Hypochaeris radicata)

(Coronopus didymus)

SCIENTIFIC NAME

(Cyprus compressus)

(Axonopris offinis)

(Digitaria spp.)

(Eleusine indica)

(Setaria viridis)

(Avena fatua)

(Bromus catharticus)

(Echinochloa crus-galli)

(Panicum dichotomiflorum)

(Amaranthus retroflexus)

(Capsella bursa-pastoris)

(Polygonum pensylvanicum)

(Oxalis spp.)

Holly Japanese - Helleri Kurogane

SPECIES

Hemlock*

Oleander

Pine

Chinese - Parsonii Juniper Common - Berkshire Effusa

> Creeping - Blue Chip, Blue Rug, Andorra

> > Singleseed - Blue Star

VARIETIES

Japanese

American

Purpurea

Gold Spot

Canada

Nootka - Aurea

Eastern Red

Dazzler

Giant - Zebrina

Gvpsophila paniculata

American - Smaragd, Rheingold

Evergreen - Golden Japanese.

Hinoki - Graciosa, Pygmaea

Douglas, Fraser, Grand, Noble

Sawara - Aurea Nana

Savin - Buffalo, Tamariscifolia Common

Ophiopogon Eastern White, Mugho, Scotch

Raphiolepis

Spruce* Statice Limonium spp.

*Not currently registered for use on Hemlock, Fir, and Blue Spruce in California.

Local conditions may affect the use of this product. Consult State Agricultural Experiment Station or Extension Service weed specialists for specific recommendations regarding local weed problems and for information about using lower dosages.

LANDSCAPE PLANTINGS: ROADSIDE, COMMERCIAL, & INDUSTRIAL SITES, Apply Quali-Pro Oxadiazon 50 WSB before weed seed germination as an over-the-top spray only on those ornamentals on this label.

USE RATES: Apply Quali-Pro Oxadiazon 50 WSB as a broadcast spray in a minimum of 50 gallons of water per acre to established ornamentals using 4 water-soluble packets per acre. Remove all existing weed growth before applying. Applications should not be made within 60 days of transplanting. Do not apply with drift-control agents or adjuvants. Sprinkle irrigate within 2 hours after applying this product; otherwise, crop injury may result.



Oxadiazon 50 WSB

Herbicide

A pre-emergent herbicide for the control of broadleaf weeds and annual grasses in turf and ornamentals.

FOR SALE TO AND USE BY PROFESSIONAL APPLICATORS ONLY. NOT FOR SALE TO OR USE BY HOMEOWNERS/CONSUMERS. NOT FOR USE ON HOME LAWNS

ACTIVE INGREDIENT	% BY WT.
Oxadiazon: 3-[2,4-dichloro-5-(1-methylethoxy)phenyl]	-
5-(1,1-dimethylethyl)-1,3,4-oxadiazol-2(3H)-one	
INERT INGREDIENTS:	50.0%
TOTAL:	100.0%

EPA Reg. No. 66222-209 EPA Est. No. 65387-AR-001

KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

For additional precautionary, handling, and use statements, see inside of this booklet

Manufactured for:

Makhteshim Agan of North America, Inc.
4515 Falls of Neuse Road, Suite 300
Raleigh. NC 27609

FIRST AID	
IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Riinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing. Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a	

poison control center or doctor or going for treatment. You may also contact Prosar at 1-877-250-9291 for emergency medical treatment information.

Note to Physician: There is no specific antidote. Appropriate supportive and symptomatic treatment as indicated by the patient's condition is recommended.

EPA 011008/Trans 010410/Notif 030510/Rev D

14158