

0-0-12

A True Solution Plant Activator with PAC* Chelation Technology

Guaranteed Analysis:

 $\begin{array}{lll} \text{Soluble Potash (K_2O)} & 12.0\% \\ \text{Calcium (Ca)} & 2.4\% \\ \text{Magnesium (Mg)} & 0.8\% \\ \end{array}$

Derived from calcium hydroxide, magnesium oxide and potassium hydroxide

For use as a fertilizer in foliar and fertigation applications to supplement K, Ca and Mg levels and to promote plant metabolism and translocation of sugars in agricultural crops.

*PAC refers to NutriAg's "Polyaldocarbosate" Chelation Technology

 $Information\ regarding\ the\ contents\ and\ levels\ of\ metals\ in\ this\ product\ is\ available\ on\ the\ internet\ at\ http://www.aapfco.org/metals.htm$

Directions for Use (Use only as Directed):

- Avoid foliar application in poor drying conditions such as very high humidity, heavy dew set or continually rainy weather.
- Do not foliar apply when ambient temperatures are expected to exceed 85°F before application completely dries on foliage
- Aerial application minimum of 5 gallons of water/acre; Dilute sprays minimum of 30 gallons of water/acre.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- Phyto B is compatible with most agricultural chemicals and nutrients. Should be applied in conjunction with tissue and/or soil analysis.
- . Consult a NutriAg representative for application timing and necessity of its use.

Suggested Application Rates:

| Crop | Foliar Rates^^ | Fertigation Rates^^ |
|---|--------------------|--------------------------|
| Forage Crops: Alfalfa, Clover, Hay Crops | | |
| Field Crops: Cereals, Corn, Cotton, Sorghum, Sugar cane | | |
| Legumes: Dry bean, Lentil, Peanut, Pea, Soybean | | |
| Root Vegetables (including, but not limited to): Beet, Carrot, Garlic, Parsnip, Radish, Rutabaga, | | The number and |
| Sugar Beet, Sweet Potato, Turnip | | frequency of |
| Leafy Vegetables (including but not limited to): Bok Choy, Celery, Chard, Cilantro, Cole Crops, | 1-2 quarts/acre/ | applications determine |
| Lettuce, Parsley, Spinach | application | application rates. |
| Other Vegetables: Artichoke, Asparagus, Cucurbits, Eggplant, Okra, Onions (bunching, bulbing), | Early and an early | America 4 Americants (|
| Peppers, Rhubarb, Snap bean, Sweet Corn, Tomato | For best results, | Apply 1-4 quarts/ |
| Berry Crops: Blueberry, Caneberry, Cranberry, Elderberry, Gooseberry, Strawberry | a minimum of | acre/application. |
| Tree Fruit (including but not limited to): Apple, Apricot, Avocado, Cherry, Nectarine, Olive, | 2 applications is | |
| Peach, Pear, Persimmon, Plum, Pomegranate, Prune, Quince | recommended. | Always apply toward |
| Other Fruit: Citrus (all), Fig, Kiwi, Melons | | the end of an irrigation |
| Nut Trees (including but not limited to): Almond, Filbert, Hazelnut, Pecan, Pistachio, Walnut | | cycle. |
| Grapes: Currant, Raisin, Table, Wine | | |
| Other Crops: Herbs, Hops, Mint, Potato, Safflower, Spearmint, Stevia, Sunflower | | |

**Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions. Lower rates are recommended for general crop maintenance or when applying multiple times per season; higher rates are recommended when applying a minimum number of applications to crops that require repeat applications, or where more severe deficiency is present. Higher volumes of water should be used for higher rates of fertilizer. For application timing charts, please consult a product manual. Consult your NutriAg representative for specific crop recommendations.

Warranty Statement: Sellers guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property anising from the use or handling of this product and accepts the product on this condition. NutriAg Ltd. warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when sed in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product such as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of NutriAg Ltd. NutriAg Ltd. will not be liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.

Caution: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with skin, eyes or clothing. In case of contact, immediately flush skin or eyes with water for at least 15 minutes.

Net Contents:

2.5 Gal. (9.46 L)

Net Weight:

25.66 lbs. (11.64 kg)

Net Weight/Gallon: 10.27 lbs. (4.66 kg)

Batch No.: XXXXXX