




US79862768D (100816) LABMC TEMIK 15 G LOCK 'N LOAD DATAPACK - 9/1/10
 COLORS: CMYK

5.75" x 3.875"

<p>RESTRICTED USE PESTICIDE DUE TO ACUTE ORAL TOXICITY and TO GROUND WATER CONTAMINATION <small>For retail sale to and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.</small></p>								
 <p>TEMIK[®] 15G <small>brand</small></p> <p>LOCK'nLOAD[®] <small>ALDICARB PESTICIDE</small></p> <p>Net Contents: 45 pounds</p> <p><i>For Control of Certain Insects, Mites, and Nematodes</i> ACTIVE INGREDIENT: Aldicarb [2-methyl-2-(methylthio)propionaldehyde Q-(methylcarbamoyl)oxime] 15% INERT INGREDIENTS: 85%</p> <p>EPA Reg. No. 264-330 EPA Est. No. 264-GA-01</p> <p>KEEP OUT OF REACH OF CHILDREN  DANGER POISON  PELIGRO</p> <p>4250701</p>	<p>Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.) FOR ADDITIONAL PRECAUTIONARY STATEMENTS: See inside booklet. For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937) For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577</p>							
	<p style="text-align: center;">FIRST AID</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 20%;">IF SWALLOWED:</td> <td> <ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person. </td> </tr> <tr> <td>IF IN EYES:</td> <td> <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice. </td> </tr> <tr> <td>IF ON SKIN OR CLOTHING:</td> <td> <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. </td> </tr> <tr> <td>IF INHALED:</td> <td> <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice. </td> </tr> </table> <p style="text-align: center;">For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577. Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person. 	IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice. 	IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. 	IF INHALED:
IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person. 							
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice. 							
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. 							
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice. 							
<p><small>Aldicarb is a N-methyl carbamate. TEMIK is a registered trademark of Bayer. Produced for: Lock 'n Load is a registered trademark of Bayer CropScience LP AMVAC Chemical Company P.O. Box 12014, 2 T.W. Alexander Drive ©2010 Bayer CropScience Research Triangle Park, North Carolina 27709 US79862768D 100816D 09/10</small></p>								

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

MEDICAL INFORMATION

Contact a physician immediately in all cases of suspected poisoning. Illness may be produced rapidly following overexposure to TEMIK brand 15G Lock 'n Load Aldicarb Pesticide. If breathing stops, establish an airway and start artificial respiration, and provide oxygen. Make certain to remove all sources of continuing contamination. Remove clothing and wash skin and hair immediately with large amounts of water. Transport the patient to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN. If poisoning is suspected in animals, contact a veterinarian.

ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN.

NOTE TO PHYSICIAN: TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is a n-methyl carbamate insecticide which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.

Gastric lavage may be used if this product has been swallowed. TEMIK brand 15G Lock 'n Load Aldicarb Pesticide poisoning may occur rapidly after ingestion and prompt removal of stomach contents is indicated.

Specific treatment consists of the administration of parenteral atropine sulfate. Caution should be exercised to prevent overatropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until the patient is fully atropinized, then intramuscularly every 30 to 60 minutes as needed to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs such as (pyridine-2-aldoxime methiodide) are NOT recommended unless organophosphate intoxication is also suggested.

To aid in confirmation of a diagnosis, urine samples must be obtained within 24 hours of exposure and immediately frozen. Analyses will be arranged by Bayer CropScience. Consultation on therapy can be obtained at all hours by calling the Bayer CropScience emergency number: 1-800-334-7577.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

FATAL IF SWALLOWED. Causes cholinesterase inhibition. May be fatal or harmful by skin or eye contact or by breathing dust. Rapidly absorbed through skin or eyes. Do not get in eyes, on skin or on clothing. Do not breathe dust. Keep away from domestic animals.

Always stand up-wind from hopper when loading.

SIGNS AND SYMPTOMS OF OVEREXPOSURE:

Salivation, Muscle tremor, Nausea, Watery eyes, Difficult breathing, Vomiting, Pinpoint eye pupils, Excessive sweating, Diarrhea, Blurred vision, Abdominal cramps, Weakness, Headache.

In severe cases, convulsions, unconsciousness, and respiratory failure may occur.

PERSONAL PROTECTIVE EQUIPMENT (PPE) FOR TEMIK BRAND 15G LOCK 'N LOAD ALDICARB PESTICIDE IN LOCK 'N LOAD AND SMARTBOX SYSTEMS.

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart.

Personal Protective Equipment (PPE):

All handlers (except loaders) must wear a minimum of a long-sleeved shirt, long pants, chemical resistant gloves made of any waterproof material, and shoes plus socks. In addition, during equipment cleaning or repair, or spill clean up, handlers must wear protective eyewear (goggles or face shield), a chemical resistant apron, and a NIOSH-approved respirator with a dust/mist filter with the MSHA/NIOSH approval number prefix TC-21C or any N, R, P or HE filter.

Engineering Controls For Closed Loading Systems:

This product is formulated into Lock 'n Load and Smartbox systems that meet the definition of a closed loading system in the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240(d)(4)]. Loaders using the closed-system packaging may wear reduced personal protective equipment provided they wear a long-sleeved shirt, long pants, and shoes plus socks and, are provided, have

(continued)

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

immediately available, and use in an emergency, such as a broken package, spill, or equipment breakdown: chemical resistant gloves made of any waterproof material, a chemical resistant apron, chemical-resistant footwear, protective eyewear (goggles or face shield), and a NIOSH-approved respirator with a dust/mist filter with the MSHA/NIOSH approval number prefix TC-21C or any N, R, P, or HE filter.

Engineering Controls For Enclosed Cab Vehicles:

Applicators using an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240(d)(5)] may wear reduced personal protective equipment provided they wear a long-sleeved shirt, long pants, and shoes plus socks and, are provided, have immediately available, and use in an emergency, such as a broken package, spill, or equipment breakdown: chemical resistant gloves made of any waterproof material, a chemical resistant apron, chemical-resistant footwear, protective eyewear (goggles or face shield), and a NIOSH-approved respirator with a dust/mist filter with the MSHA/NIOSH approval number prefix TC-21C or any N, R, P, or HE filter. Applicators must take off any PPE that was worn in the treated area before reentering the cab, and store all such PPE in a chemical-resistant container, such as a plastic bag, to prevent contamination of the inside of the cab.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

TOXIC TO FISH, BIRDS, AND OTHER WILDLIFE: This pesticide is extremely toxic

to birds and other wildlife. Birds feeding on exposed granules may be killed. Cover or immediately soil incorporate granules spilled during loading, at row ends, or elsewhere to ensure the granules are completely covered with soil.

This pesticide is toxic to fish and aquatic invertebrates. Run-off from treated areas may be hazardous to fish in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is potentially toxic to honey bees through translocated residues in pollen and nectar if application is made during bloom.

Aldicarb is known to leach through soil into ground water under certain conditions. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination. Apply this product only as specified on this label. Read the use directions and the appended Environmental Precautions and Soil Type Restriction Tables prior to making applications. If there are any questions, contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937).

DECOMPOSITION AND MOVEMENT IN SOIL: This product is readily decomposed into harmless residues under most use conditions. However, a combination of permeable and acidic soil conditions, moderate to heavy irrigation and/or rainfall, use of 20 or more pounds per acre, and soil temperature below 50°F (10°C) at application time tend to reduce degradation and promote movement of residues to ground water. If the above describes your local use conditions and ground water in your area is used for drinking, do not use this product without first contacting Bayer CropScience at 1-866-99BAYER (1-866-992-2937).

NOTICE: Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species.

This Act protects Attwater's Greater Prairie Chicken in the Texas counties of Aransas, Austin, Brazoria, Colorado, Galveston, Goliad, Harris, Refugio, and Victoria.

Prior to making applications in these counties, the user must determine that this species is not located in or immediately adjacent to the area to be treated. If the user is in doubt whether or not the above named endangered species may be affected, he should contact either the regional U.S. Fish & Wildlife Service office (Endangered Species Specialist) or personnel of the State Fish and Game office.

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

Exception: If the product is soil-incorporated or soil-injected, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. Once the REI has expired, do not allow persons to come into direct contact with treated soil wet as the result of the initial irrigation or rainfall after treatment unless they are wearing the PPE specified below for early entry. PPE required for early entry to treated areas that is permitted under the Worker Protections Standard and that involves contact with anything that has been treated, such as plants, soil, or water is coveralls worn over long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, chemical-resistant footwear plus socks, and protective eyewear.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

STORAGE AND DISPOSAL

STORAGE

Store unused TEMIK brand 15G Lock 'n Load Aldicarb Pesticide in original container only, in secure, well ventilated clean dry area out of reach of children and animals. Do not store in areas where temperature averages 115°F (46°C) or greater. Do not store in or around the home or home garden.

PESTICIDE DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL

FOR LOCK 'N LOAD AND SMARTBOX PACKAGING ONLY Refillable container. Refill this container with TEMIK brand 15G Lock 'n Load Aldicarb Pesticide only. Do not reuse this container for any other purpose. Return empty Lock 'n Load and Smartbox containers to point of purchase.

IN CASE OF TRANSPORTATION OR WAREHOUSE EMERGENCY INVOLVING A SPILL, FIRE, OR EXPOSURE, CALL 1-800-334-7577 TWENTY-FOUR HOURS A DAY IN THE USA.

INFORMATION

READ ENTIRE LABEL BEFORE USING THIS PRODUCT.

TEMIK brand 15G Lock 'n Load Aldicarb Pesticide controls listed insects, mites, and nematodes. When applied into moist soil at planting and/or postemergence the active ingredient aldicarb is rapidly absorbed by roots and translocated to all parts of the plant. Rainfall or irrigation soon after application will ensure prompt uptake of aldicarb; however, if irrigation is necessary care should be taken not to over irrigate to reduce the potential for residues leaching to ground water. Control often lasts more than six weeks varying with growing conditions, rate of use, and pests. Crop yields are usually increased with treatments of TEMIK brand 15G Lock 'n Load Aldicarb Pesticide.

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

RESTRICTIONS

Use TEMIK brand 15G Lock 'n Load Aldicarb Pesticide only in accordance with label directions, and restrictions. **DO NOT USE ON ANY CROP NOT LISTED ON THIS LABEL OR SUPPLEMENTAL LABELING.** Use higher rates on heavy organic or clay soils. Do not exceed the maximum label rate.

DO NOT APPLY TEMIK BRAND 15G LOCK 'N LOAD ALDICARB PESTICIDE TO ANY CROP AFTER AUGUST 31, 2018.

Application may only be made using motorized ground application equipment. Application using aircraft, backpack spreaders, or push-type spreaders is prohibited. Additional restrictions on application equipment pertain to use on sweet potatoes.

Make side-dress applications close enough to plants to allow good uptake by the roots without injury to the plants from root pruning. In irrigated areas, follow application with irrigation within one week. If alternate furrows are irrigated after side-dress application, TEMIK brand 15G Lock 'n Load Aldicarb Pesticide and water must be on the same side of the plant row.

Calibrate and adjust ground application equipment to insure proper rate and accurate placement. Clean application equipment thoroughly after use. For any left over material, see instructions for STORAGE AND DISPOSAL in this booklet.

Deep disk any spills at row ends immediately to prevent birds from feeding on exposed granules.

GROUND WATER RESTRICTIONS

- Observe Environmental Hazard statements regarding Decomposition and Movement in Soil, and carefully follow Directions For Use.
- In fields having soils with less than 15% field moisture holding capacity, special care must be taken not to over-irrigate, since over-irrigation promotes the leaching of chemicals.
- **Do not apply within 50 feet of any drinking water well. More restrictive setbacks may apply. See Directions For Use and the *Environmental Precautions and Soil Type Restriction Tables*.**
- Do not wash, load or empty application equipment near any well, as this practice is a potential source of ground water contamination.
- Do not apply this product in the states of Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, Vermont or Wisconsin, or in Del Norte or Humboldt counties in California or in Curry County, Oregon.

- For State Specific Ground Water Restrictions, see Directions For Use and the *Environmental Precautions and Soil Type Restriction Tables*.
- Apply only between March 1 and September 1 when used in California.

ROTATIONAL CROPS RESTRICTION

Do not plant any crop not listed on this label in soil treated with TEMIK brand 15G Lock 'n Load Aldicarb Pesticide within 10 months after the last application with the following exceptions:

Six-month plantback restriction

- Do not plant wheat or barley within 6 months after last application.
- Do not plant bulb crops (such as onions or garlic) within 6 months after the last application.
- Do not plant brassica crops (such as broccoli or cabbage) within 6 months after the last application.

Eight-month plantback restriction

- Do not plant corn within 8 months of last application.
- Do not plant melons within 8 months after last application.
- Do not plant other cucurbits (such as cucumbers and squash) within 8 months after last application.
- Do not plant fruiting vegetables (such as tomatoes or eggplant) within 8 months after the last application.

MEAT AND MILK

- Do not allow livestock to graze in treated areas before harvest.

COMPATIBILITY

Pesticidal activity of TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is not affected by normal applications of fertilizers or other pesticides. Its effectiveness may be reduced or lost if applied with alkaline materials such as lime. To minimize potential exposure hazards, do not mix TEMIK brand 15G Lock 'n Load Aldicarb Pesticide with other materials before application.

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

DIRECTIONS FOR USE

To provide maximum performance and to minimize hazard to birds, granules must be placed into bottom of furrow and immediately covered with soil by mechanical means. For all applications, cover or immediately soil incorporate granules spilled during loading, at row ends, or elsewhere to ensure the granules are completely covered with soil. When a range of rates are specified, use the higher rate if pest infestations are expected to be severe.

COTTON				
Crop & Time Of Application	Pests Controlled	Pounds/ Acre	EXAMPLE Ounces/ 1000 Feet of Row	Application Instructions
<p>COTTON At planting</p> <p>In the states of AL, FL, GA and SC, if a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>	Aphids, Thrips	3.5 to 5	<p>40' row spacing 4.5 to 6</p> <p>(Except Texas, Oklahoma, and New Mexico)</p>	<p>Apply granules in the seed furrow and immediately cover with soil by mechanical means.</p> <p>If seeds and TEMIK brand 15G Lock 'n Load Aldicarb Pesticide are hill-dropped, TEMIK brand 15G Lock 'n Load Aldicarb Pesticide rates may be reduced by one-half.</p>
		2 to 5	2.5 to 6 (Texas, Oklahoma, and New Mexico Only)	
	Fleahoppers, Leafminers, Mites, Overwintering boll weevil (adults feeding on foliage), Plant bugs, including Lygus	4 to 7	5 to 8.5	
	Nematodes (Suppression)	4 to 7	5 to 8.5 (California Only)	Apply in seed furrow and cover with soil.
	Nematodes	3.5 to 7	4.5 to 8.5 (Except California)	Apply granules in seed furrow and immediately cover with soil by mechanical means. OR Apply granules in a 4 to 6-inch band (T-Band) over open seed furrow and immediately cover with soil by mechanical means.

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

COTTON (continued)

Crop & Time Of Application	Pests Controlled	Pounds/Acre	EXAMPLE Ounces/1000 Feet of Row	Application Instructions
<p>Side Dress Application only or Split Application Make side dress applications from 3 weeks after planting through first squaring. Side Dress Application only In the states of AL, FL, GA and SC, if a Side Dress application only is planned and if a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p> <p>Split Application In the states of AL, FL, GA and SC, if an At planting and Side Dress application is planned and if a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1000 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>	Suppression of Leafhoppers, Fleahoppers, Mites, Lygus and Nematodes	5	6	Side dress granules in a furrow that is 6 to 10 inches to one or both sides of plant row to a depth of 2 to 3 inches. Adjust applications to minimize root pruning.

RESTRICTIONS FOR COTTON

- The maximum single at-plant application rate is 7 pounds per acre. See application restrictions on vulnerable soils.
- The maximum single side-dress application rate is 5 pounds per acre. See application restrictions on vulnerable soils.
- Do not exceed a total of 12 pounds per acre per year for all applications to cotton. See application restrictions on vulnerable soils where a combination of an At-Planting and Side Dress application is planned.
- Apply only between March 1 and September 1 when used in California.
- Do not make more than one at-planting application and one postemergence application per crop per year.
- Make side-dress applications close enough to plants to allow good uptake by the roots without injury to the plants from root pruning.
- Do not apply within 90 days of harvest.
- Do not feed cotton forage to livestock or allow livestock to graze in treated area.
- Immediately deep-disk any spills at row ends or elsewhere to ensure the granules are covered with a layer of soil.

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

DRY BEANS				
Crop & Time Of Application	Pests Controlled	Pounds/ Acre	EXAMPLE Ounces/ 1000 Feet of Row	Application Instructions
DRY BEANS At planting FOR USE ONLY IN: Colorado, Oregon, Washington, Idaho, and Michigan.	Seedcorn maggot	3.5 to 5	36 inch row spacing 4.0 to 5.5	Apply granules in seed furrow and immediately cover with soil by mechanical means. OR
	Aphids	5 to 7	5.5 to 7.5	Apply in a furrow that is 2 to 3 inches to side of seed row and 2 to 3 inches deep and immediately cover with soil by mechanical means. OR
	Leafhoppers	7 to 14	7.5 to 15	Granules can be placed in seed furrow if the rate does not exceed 5.5 ounces per 1000 feet of row (5 pounds per acre on 36 inch rows)
	Mexican bean beetle Mites			
	Nematodes	7 to 14	7.5 to 15	Apply granules in a 4 to 6 inch band and immediately cover with soil by mechanical means. Plant into treated zone. OR Where furrow irrigation is used, apply granules 3 to 4 inches deep and 3 inches from seed row on the water furrow side.

RESTRICTIONS FOR DRY BEANS

- For use only in Colorado, Oregon, Washington, Idaho, and Michigan.
- Do not exceed a total of 14 pounds per acre per season.
- Treatments in excess of 5.5 ounces per 1000 feet of row (5 pounds per acre on 36 inch rows) made directly to the seed furrow may delay plant emergence and reduce plant stand.
- Do not make more than one application per crop per year.
- Do not harvest within 90 days of treatment.
- Do not feed green forage, hay or straw to livestock.
- Do not allow livestock to graze in treated areas before harvest.
- Do not use green pods as food for humans.
- Immediately deep-disk any spills at row ends or elsewhere to ensure the granules are covered with a layer of soil.

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

PEANUTS

Crop & Time Of Application	Pests Controlled	Pounds/ Acre	EXAMPLE Ounces/ 1000 Feet of Row	Application Instructions
PEANUTS At planting In the states of AL, FL, GA and SC, if a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.	Thrips	7	36" row spacing 7.5	Apply granules in seed furrow and immediately cover with soil by mechanical means. In Southwest United States use high rate only.
	Nematode suppression	7	7.5	Apply granules in a 4 to 6 inch band over open seed furrow (T-band) and immediately cover with soil by mechanical means. OR Apply granules in a 6 to 12 inch band and immediately cover with soil by mechanical means to a depth of 2 to 4 inches. Plant seed into treated zone.
Split Application (Alabama, Florida, Georgia, North Carolina, Oklahoma, Texas and Virginia Only) At planting and/or Post-emergence (pegging application) at or just prior to peg initiation but no later than 40 days after emergence and prior to last cultivation. In the states of AL, FL, and GA, if an At planting and/or a Pegging application is planned and if a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1100 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.	Nematodes	7 At-planting followed by 10 Post-emergence	7.5 At-planting followed by 11 Post-emergence	At-Planting: Apply granules in a 4 to 6 inch band over open seed furrow (T-band) and immediately cover with soil by mechanical means. OR Apply granules in a 6 to 12 inch band and immediately cover with soil by mechanical means to a depth of 2 to 4 inches. Plant seed into treated zone. Post-Emergence: Apply granules in a band 12 to 18 inches wide on the row and into the plant canopy. Ensure that plant foliage is dry prior to application. Dislodge granules from foliage by suitable means that will not damage the plant. Irrigate immediately after application and complete within 24 hours unless rainfall is received.

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

PEANUTS (continued)

RESTRICTIONS FOR PEANUTS

- Do not make more than one application per crop per year in states other than Alabama, Florida, Georgia, North Carolina, Oklahoma, Texas and Virginia.
- The maximum single at-plant application rate is 7 pounds per acre. See application restrictions on vulnerable soils.
- The maximum single Pegging application rate is 10 pounds per acre. See application restrictions on vulnerable soils.
- Do not exceed a total of 17 pounds per acre per year for all applications to peanuts. See application restrictions on vulnerable soils where a combination of an At-Planting and pegging application is planned.
- Post-emergence applications are permitted only in fields where overhead irrigation is available.
- Do not make the split application to Spanish peanuts or other short season varieties (a minimum of 90 days is required between pegging applications and harvest).
- Do not harvest within 90 days of application.
- Do not hog-off treated fields.
- Do not allow livestock to graze in treated areas before harvest.
- Do not feed hay or vines to livestock.
- Immediately deep-disk any spills at row ends or elsewhere to ensure the granules are covered with a layer of soil.

SOYBEANS

Crop & Time Of Application	Pests Controlled	Pounds/ Acre	EXAMPLE Ounces/ 1000 Feet of Row	Application Instructions
SOYBEANS At planting FOR USE ONLY IN: Georgia, North Carolina, South Carolina, and Virginia In the states of GA and SC, if a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.	Mexican bean beetle Thrips Suppression of three cornered alfalfa hopper	5 to 7	30" <u>row spacing</u> 4.6 to 6.4	Apply granules in seed furrow and immediately cover with soil by mechanical means. Granules may be applied in seed furrow if rate does not exceed 5.5 ounces per 1000 feet of row OR If rate exceeds 5.5 ounces per 1000 feet of row (6 pounds per acre on 30 inch rows), apply a 4 to 6 inch band over open seed furrow and immediately cover with soil by mechanical means.

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOYBEANS (continued)

Crop & Time Of Application	Pests Controlled	Pounds/Acre	EXAMPLE Ounces/1000 Feet of Row	Application Instructions
<p>SOYBEANS At planting FOR USE ONLY IN: Georgia, North Carolina, South Carolina, and Virginia In the states of GA and SC, if a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>	<p>Nematodes (suppression)</p>	<p>7</p>	<p>6.4</p>	<p>Apply a 4 to 6 inch band (T-Band) over open seed furrow and immediately cover with soil by mechanical means.</p>

RESTRICTIONS FOR SOYBEANS

- For use only in Georgia, North Carolina, South Carolina, and Virginia.
- Do not make more than one application per crop per year.
- Do not harvest within 90 days of treatment.
- Do not allow livestock to graze in treated areas before harvest.
- Do not feed green forage, hay, or straw to livestock.
- Treatments in excess of 5.5 oz per 1000 feet of row (6 pounds per acre on 30 inch rows) made directly to the seed furrow may delay plant emergence and reduce plant stand.
- Immediately deep-disk any spills at row ends or elsewhere to ensure the granules are covered with a layer of soil.

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SUGAR BEETS				
Crop & Time Of Application	Pests Controlled	Pounds/Acre	EXAMPLE Ounces/1000 Feet of Row	Application Instructions
SUGAR BEETS At planting or within one week before planting. FOR USE ONLY IN: California, Colorado, Idaho, Montana, Nebraska, Oregon, Washington and Wyoming.	Nematodes	27 to 33 (Except California)	22" row spacing 18 to 22	Apply granules in a 4 to 6 inch band and immediately cover with soil by mechanical means. Plant seed into or above treated zone; OR where furrow irrigation is employed for seed germination, drill granules 3 to 4 inches deep and 3 inches from seed row on water furrow side.
		14 (California Only)	9.5	
At planting	Aphids	7 to 14	4.5 to 9.5	Drill granules 1 to 3 inches below seedline. Granules can be placed in seed furrow if rate does not exceed 7 pounds per acre. Repeat applications will be required for continued protection against virus vectors (aphids and leafhoppers)
	Leafminers Leafhoppers	14 to 20 (Except California)	9.5 to 13.5	
		14 (California Only)	9.5	
Sugar beet root maggot	7 to 14	4.5 to 9.5	Apply granules in a 2 to 3 inch band over seed row and immediately cover with soil by mechanical means. OR where furrow irrigation is employed for seed germination, drill granules 2 inches deep and 2 inches from seed row on water furrow side. For the 7 lb. rate, apply granules in a 1 to 2-inch band in front of the press wheel as the furrow is closing.	

(continued)

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SUGAR BEETS (continued)

Crop & Time Of Application	Pests Controlled	Pounds/Acre	EXAMPLE Ounces/1000 Feet of Row	Application Instructions
At planting plus postemergence (split applications)	Nematodes	14 to 20 at planting and 14 to 20 at postemergence (Except California)	9.5 to 13.5 at planting and 9.5 to 13.5 at postemergence	AT PLANTING: Apply granules in a 4 to 6-inch band and immediately cover with soil by mechanical means. Plant seed into or above treated zone; OR where furrow irrigation is employed for seed germination, drill granules 3 to 4 inches deep and 3 inches from seed row on water furrow side. POSTEMERGENCE: Apply granules to both sides of plant row and immediately cover with soil by mechanical means. OR where furrow irrigation is employed side-dress granules 4 to 8 inches to water furrow side of plant row at furrow depth. Irrigate soon after application. Apply within 60 days after planting.
		14 at planting and 14 at postemergence (California Only)	9.5 at planting and 9.5 at postemergence	
Postemergence Do not make any postemergence application if 27 to 33 lbs./acre were applied at planting or one week before planting. Do not make more than one at planting application and two postemergence applications per crop.	Sugar beet root maggot	7 to 14	4.5 to 9.5	Apply granules to both sides of plant row and immediately work into the soil or cover with soil by mechanical means OR where furrow irrigation is employed side-dress granules 4 to 8 inches to water furrow side of plant row at furrow depth. Irrigate soon after application. Apply within 60 days after planting. OR Apply as above. A repeat application may be required for continued protection against virus vectors (aphids, leafhoppers). Apply within 60 days after planting.
	Aphids	7 to 14	4.5 to 9.5	
	Leafminers Leafhoppers	14 to 20 (Except California)	9.5 to 13.5	
		14 (California Only)	9.5	

(continued)

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SUGAR BEETS (continued)

Crop & Time Of Application	Pests Controlled	Pounds/Acre	EXAMPLE Ounces/1000 Feet of Row	Application Instructions
Postemergence Do not make any postemergence application if 27 to 33 lbs./acre were applied at planting or one week before planting. Do not make more than one at planting application and two postemergence applications per crop.	Nematodes	27 (Except California)	18	Apply as above. Apply within 60 days after planting. In California, apply within 30 days after planting.
		14 (California Only)	9.5	

RESTRICTIONS FOR SUGAR BEETS

- For use only in California, Colorado, Idaho, Montana, Nebraska, Oregon, Washington and Wyoming.
- Apply only between March 1 and September 1 when used in California.
- Do not exceed a total of 28 pounds per acre per year in California and 33 pounds per acre per year in other states for all applications to sugar beets.
- Do not make more than one at planting application and two postemergence applications per crop per year.
- Do not apply within 90 days of harvest.
- If tops are to be fed to livestock, do not apply within 120 days of harvest.
- Do not use tops as food for humans.
- Treatments in excess of 4.5 ounces per 1000 ft of row (7 pounds per acre on 22 inch rows) made directly in the seed furrow may delay plant emergence and reduce plant stand.
- Immediately deep-disk any spills at row ends or elsewhere to ensure the granules are covered with a layer of soil.

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SWEET POTATOES

Crop & Time Of Application	Pests Controlled	Pounds/Acre	EXAMPLE Ounces/1000 Feet of Row	Application Instructions
SWEET POTATOES Pre-plant or at planting FOR USE ONLY IN: Louisiana and Mississippi	Nematodes	10 to 20	<u>48" row spacing</u> 15 to 30	Apply granules in a 12-inch band over open furrow or soil surface and cover immediately during bed forming by mechanically hilling up 8 to 10 inches. Place transplant in center of treated zone.

REQUIRED APPLICATION EQUIPMENT FOR SWEET POTATOES

All applications must be made with Positive Displacement Applicators such as Horstine Microband® or Gandy PDM® or with other Positive Displacement Applicators.

RESTRICTIONS FOR SWEET POTATOES

- For use only in Louisiana and Mississippi.
- Maximum single application rate is 20 pounds per acre per year.
- Do not make more than one application per crop per year.
- Do not harvest within 120 days of application.
- Do not feed sweet potato vines to livestock.
- Immediately deep-disk any spills at row ends or elsewhere to ensure the granules are covered with a layer of soil.

PLANT GROWTH STIMULATION: In the absence of recognizable target pests and under certain growing conditions, stimulation of plant growth by TEMIK brand 15G Lock 'n Load Aldicarb Pesticide on certain crops has been demonstrated under laboratory, greenhouse, and field conditions. Plants may be visibly taller, greener, and denser. Faster grow-off and an increase in fruiting rate and size often are results of the growth enhancement by TEMIK brand 15G Lock 'n Load Aldicarb Pesticide. However, such effects do not occur with all crops or under all conditions. Therefore, this product should not be used solely as a plant growth regulator, and its effect as one should be considered a side benefit which may or may not occur as a consequence of its use.

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

TEMIK BRAND 15G LOCK 'N LOAD ALDICARB PESTICIDE (GYPSUM) CALIBRATION GUIDE

APPROXIMATE APPLICATOR SETTINGS* FOR TEMIK BRAND 15G LOCK 'N LOAD ALDICARB PESTICIDE (GYPSUM) AT 4, 5 & 6 MPH

Refer to equipment manufacture literature for additional calibration instructions.

NOTE: This calibration chart is applicable only to TEMIK brand 15G Lock 'n Load Aldicarb Pesticide in this container which is formulated for use on a gypsum carrier. All rates are APPROXIMATIONS and must be confirmed by using a calibration tube (See Calibration Chart provided with tube). Calibration tubes are available from your TEMIK brand 15G Lock 'n Load Aldicarb Pesticide supplier.

TEMIK BRAND 15G LOCK 'N LOAD ALDICARB PESTICIDE (GYPSUM) CALIBRATION GUIDE

POUNDS TEMIK BRAND 15G LOCK 'N LOAD ALDICARB PESTICIDE PER ACRE FOR VARIOUS ROW SPACINGS						TYPE OF GRANULAR APPLICATOR											
						GANDY GAUGE SETTING			JOHN DEERE MAX EMERGE GAUGE SETTING**			JOHN DEERE MAX EMERGE II GAUGE SETTING			CASE-IH SERIES 800/900 GAUGE SETTING****		
ROW SPACING						MPH			MPH			MPH			MPH		
22"	34"	36"	40"	48"		4	5	6	4	5	6	4	5	6	4	5	6
5.5	3.5	3.3	3	2.5		12	13	14	6***	7***	8	8	9	11	1/6.0	1/6.5	1/7.5
7.3	4.7	4.4	4	3.3		13	15	16	7***	9	10	10	12	13	1/7.0	1/7.5	1/8.5
9.1	5.9	5.5	5	4.2		14	16	18	9	11	13	12	14	15	1/8.0	1/8.5	1/9.5
10.9	7.1	6.7	6	5.0		16	18	19	11	13	15	13	15	18	1/9.0	1/9.5	2/1.0
12.7	8.2	7.8	7	5.8		17	19	21	12	15	18	15	18	21	1/9.5	2/0.5	2/2.0
14.5	9.4	8.9	8	6.7		18	20	22	14	17	20	16	20	23	2/0.0	2/2.0	2/3.5
16.4	10.6	10.0	9	7.5		19	21	23	16	19	23	18	22	27	2/1.0	2/3.0	2/5.0
18.2	11.8	11.1	10	8.3		20	22	24	17	21	25	20	25	29	2/2.0	2/4.0	2/6.5
21.8	14.1	13.3	12	10.0		22	24	26	20	25	28	23	29	35	2/3.5	2/7.0	3/0.0
25.5	16.5	15.6	14	11.7		23	26	29	24	28	30	27	35	41	2/5.5	2/8.5	3/4.5
	18.8	17.8	16	13.3		25	28	31	26	29	32	31	39	45	2/7.5	3/2.5	3/8.0
	21.2	20.0	18	15.0		26	30	33	28	31	33	36	44	48	3/0.5	3/6.0	4/2.0
	23.5	22.2	20	16.7		28	31	34	30	32	35	40	46	50	3/2.5	3/9.0	4/5.5
			24	20.0		31	34	37	32	34	37	45	49	53	3/7.5	4/6.0	5/3.0
			27	22.5		33	36	40	33	36	38	48	52	56	4/2.5	5/0.0	5/9.5

* All settings for one hopper box outlet per row **Odd Numbered Housing (0 – 45)
 *** For John Deere gauge setting below 8 there will be great variations in flow rate. A gauge setting of 6 is the lowest recommended setting.
 **** Gate/Dial

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

HORSTINE FARMERY MICROBAND* (Using one 3/16" rotor per outlet, part #094006)											
POUNDS TEMIK BRAND 15G LOCK 'N LOAD ALDICARB PESTICIDE (GYPSUM) PER ACRE FOR VARIOUS ROW SPACINGS											
ROW SPACING	SPROCKET** COMBINATION					DRIVER/DRIVEN					
	19-34	22-34	25-34	28-34	25-28	22-22	28-25	34-28	34-25	34-22	34-19
22"	5.50	6.40	7.30	8.20	8.90	10.00	11.20	12.10	13.60	15.40	17.80
30"	4.00	4.60	5.30	5.90	6.50	7.30	8.10	8.80	9.90	11.20	13.00
36"	3.30	3.90	4.40	5.00	5.40	6.10	6.80	7.30	8.20	9.40	10.90
40"	3.00	3.50	4.00	4.50	4.80	5.50	6.10	6.60	7.40	8.40	9.80
48"	2.50	2.90	3.30	3.70	4.00	4.60	5.10	5.50	6.20	7.10	8.20
RATIO:	0.550	0.640	0.730	0.820	0.890	1.000	1.120	1.210	1.360	1.545	1.789

- The Microband is based on positive metering. Settings do not need to be changed with different tractor speeds.
- All settings for one hopper box outlet per row.
- For alternative dose rates, different width rotors can be fitted. For example, one 3/8" wide rotor will deliver twice the rates indicated in the above table.
- * Calibrations based upon use of a Horstine Farmery or Canaan 29.5 inch diameter Landwheel drive.
- ** Sprocket numbers, example: 19 - 34, indicate number of teeth on the DRIVER and DRIVEN Sprockets.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather

conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. To the extent allowed by applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT ALLOWED BY APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT ALLOWED BY APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT ALLOWED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

RESTRICTED USE PESTICIDE

DUE TO ACUTE ORAL TOXICITY and TO GROUND WATER CONTAMINATION

For retail sale to and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.

TEMIK® brand 15G Lock 'n Load Aldicarb Pesticide

EPA Reg. No. 264-330

**Environmental Precautions and
Soil Type Restriction Tables**



Refer to the container label for additional use precautions and directions.

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

TEMIK® Brand 15G Aldicarb Pesticide

EPA REG. No. 264-330

**ENVIRONMENTAL PRECAUTIONS
GENERAL USE RESTRICTIONS**

**AGRICULTURAL CHEMICALS HAVE THE POTENTIAL TO MOVE INTO SHALLOW GROUND WATER.
THE FOLLOWING RESTRICTIONS HAVE BEEN DEVELOPED TO PROTECT DRINKING WATER SUPPLIES.
DO NOT APPLY WITHIN 50 FEET OF ANY DRINKING WATER WELL TO MINIMIZE CONTAMINATION BY SURFACE RUNOFF.
MORE STRINGENT RESTRICTIONS MAY BE REQUIRED, AS DISCUSSED BELOW.**

Do not wash, load, or empty application equipment near any well, as this practice is a potential source of ground water contamination. In fields having soils with less than 15% field moisture holding capacity, special care must be taken not to over-irrigate, since substantial over-irrigation promotes the leaching of chemicals.
Contact your state pesticide regulatory authority for further information on state requirements for the use of this product.

Some states have or may be developing more restrictive regulations regarding the use and application of TEMIK brand 15G Lock 'n Load Aldicarb Pesticide. Follow all state regulations restricting the use and application of this product, including limitations on applications near drinking water sources. In all cases, the more restrictive requirements must be followed. It is the responsibility of the applicator to document the construction of wells claimed not to be shallow.

STATE SPECIFIC GROUND WATER LIMITATIONS

Do not apply this product in the states of Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, Vermont or Wisconsin, or in Del Norte or Humboldt counties in California or in Curry County, Oregon.

FOLLOW THE LISTED "ADDITIONAL RESTRICTIONS" if the following conditions are present:

STATE	SOILS FOR WHICH RESTRICTIONS APPLY	ADDITIONAL RESTRICTIONS
CO, DE, KS, KY, LA, MD, MO, MS, NC, TN, VA, WV	Loamy sand or sand surface soils and subsoils with an average organic matter in the upper 12 inches of less than 2% by weight. See <i>SOIL TYPE RESTRICTION TABLES</i> for specific soil types.	If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 300 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.
IA, IL, IN, MI, MN, MT, ND, NE, OH, SD, WY	Sandy loam, loamy sand, or sand surface soils, and loamy sand or sand subsoils, with an average organic matter in the upper 12 inches of less than 2% by weight. See <i>SOIL TYPE RESTRICTION TABLES</i> for specific soil types.	If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 500 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL RESTRICTION TABLE (continued)

STATE	SOILS FOR WHICH RESTRICTIONS APPLY	ADDITIONAL RESTRICTIONS
AL, GA, SC	<p>Vulnerable Cotton and Peanut Soils in AL, GA and SC and Vulnerable Soybean Soils in GA and SC.</p> <p>Loamy sand or sand surface soils and subsoils with an average organic matter in the upper 12 inches of less than 2% by weight.</p> <p>See SOIL TYPE RESTRICTION TABLES for specific soil types.</p>	<p>At planting application only to cotton, peanuts, or soybeans</p> <p>If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to cotton, peanuts or soybeans as an At-plant application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p> <p>Side Dress Application Only or Split Application to cotton</p> <p>If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to cotton as a Side Dress application only and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p> <p>If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to cotton as an At-plant application and a Side Dress application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1000 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p> <p>At-planting and/or Post emergence application to peanuts</p> <p>If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to peanut as an At-plant application and/or a Post emergence (Pegging) application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1100 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. Do not make a post emergence application to peanuts in South Carolina.</p>

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL RESTRICTION TABLE (continued)

STATE	SOILS FOR WHICH RESTRICTIONS APPLY	ADDITIONAL RESTRICTIONS
FL	<p>Vulnerable Peanut and Cotton Soils</p> <p>Loamy sand or sand surface soils and subsoils with an average organic matter in the upper 12 inches of less than 2% by weight. Any soil classified with a soil hydrologic group of C or D is not considered a restricted soil regardless of other properties.</p>	<p>At planting application only</p> <p>If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied At-planting to peanuts or cotton and a vulnerable soil is present (see soils listed below), and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table.</p> <p>Side Dress Application Only or Split Application to cotton</p> <p>If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to cotton as a Side Dress application only and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 700 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p> <p>If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to cotton as an At-plant application and a Side Dress application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1000 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p> <p>At-planting and/or Post emergence application to peanuts</p> <p>If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to peanut as an At-plant application and/or a Post emergence (Pegging) application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within 1100 feet of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.</p>

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL RESTRICTION TABLE (continued)

STATE	SOILS FOR WHICH RESTRICTIONS APPLY	ADDITIONAL RESTRICTIONS																														
FL	Vulnerable Peanut and Cotton Soils Loamy sand or sand surface soils and subsoils with an average organic matter in the upper 12 inches of less than 2% by weight. Any soil classified with a soil hydrologic group of C or D is not considered a restricted soil regardless of other properties.	<p>Additional Information</p> <p>In Florida, it is the responsibility of the applicator to document the construction of wells claimed not to be shallow. This must consist of (a) a copy of the well completion report issued by the appropriate water management district; or (b) a statement certified as to accuracy by a licensed well contractor. The U.S.D.A. Natural Resources Conservation Service which serves your county can tell you if the soils on your farm fall within the following types of vulnerable soils. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. See Section 5E-2.028, Florida Administrative Code (F.A.C.), for additional restrictions.</p> <table border="0" data-bbox="884 760 1541 878"> <tr> <td>Adamsville</td> <td>Candler</td> <td>Kershaw</td> <td>Orsino</td> <td>St. Lucie</td> </tr> <tr> <td>Alaga</td> <td>Eglin</td> <td>Klej</td> <td>Ortega</td> <td>Tavares</td> </tr> <tr> <td>Alpin</td> <td>Florahome</td> <td>Lake</td> <td>Paola</td> <td>Valdosta</td> </tr> <tr> <td>Archbold</td> <td>Fort Meade</td> <td>Lakeland</td> <td>Penney</td> <td></td> </tr> <tr> <td>Astatula</td> <td>Foxworth</td> <td>Neilhurst</td> <td>Quartzsammets</td> <td></td> </tr> <tr> <td>Bigbee</td> <td>Gainesville</td> <td>Orlando</td> <td>Satellite</td> <td></td> </tr> </table>	Adamsville	Candler	Kershaw	Orsino	St. Lucie	Alaga	Eglin	Klej	Ortega	Tavares	Alpin	Florahome	Lake	Paola	Valdosta	Archbold	Fort Meade	Lakeland	Penney		Astatula	Foxworth	Neilhurst	Quartzsammets		Bigbee	Gainesville	Orlando	Satellite	
Adamsville	Candler	Kershaw	Orsino	St. Lucie																												
Alaga	Eglin	Klej	Ortega	Tavares																												
Alpin	Florahome	Lake	Paola	Valdosta																												
Archbold	Fort Meade	Lakeland	Penney																													
Astatula	Foxworth	Neilhurst	Quartzsammets																													
Bigbee	Gainesville	Orlando	Satellite																													
FL	Other Soils, All Crops	State regulations require that TEMIK brand 15G Lock 'n Load Aldicarb Pesticide not be applied within 300 feet of any drinking water well. See Section 5E-2.028, F.A.C., and additional restrictions for peanut and cotton above.																														

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL TYPE RESTRICTION TABLE

FOLLOW THE LISTED ADDITIONAL RESTRICTIONS IF THE SOIL TYPES ARE PRESENT IN YOUR STATE:

ALABAMA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. **At planting application only:** If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to **cotton or peanuts** as an At-plant application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **700 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. **Side Dress Application Only:** If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to **cotton** as a Side Dress application only and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **700 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. **Split Application:** If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to **cotton** as an At-plant application and a Side Dress application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **1000 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. **At-planting and/or Post emergence application to peanuts:** If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to peanut as an At-plant application and/or a Post emergence (Pegging) application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **1100 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

alaga	bigbee	chiple	duckston	fripp	lakeland	osier	scranton
alpin	blanton	chipola	eustis	garcon	latonia	pactolus	tarboro
americus	bruno	corolla	fiomaton	gorgas	leon	plummer	
bassfield	buncombe	crevasse	foxworth	jones	nugent	saffell	

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL TYPE RESTRICTION TABLE (continued)

COLORADO

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

bangston	columbine	eghelm	ipage	lonetree	osgood	schamber	tomichi
bankard	comad	ellicott	ironsprings	luning	ouray	schooner	trail
barcus	corlena	els	juget	mathis	pando	sheppard	tricera
baroid	corlett	elsmere	julesburg	maybell	patna	siebert	tulloch
batterson	costilla	falcon	kandaly	medano	peetz	southace	valent
bijou	cotopaxi	farb	kassler	mespun	pescar	space city	valentine
blakeland	coyot	farisita	kerber	mido	pineguest	sphinx	wigton
boel	crestman	gilcrest	kettle	mirror lake	platte	spool	willwood
brad	crustown	gracot	kippen	moenkopie	redcreek	stapleton	yetull
breece	dailey	gretdivid	laird	moosed	resort	stecum	zeona
bresser	dix	grieves	laney	mosca	roswell	tassel	
canlodore	dunday	grimm	las animas	munjor	ruedloff	thoroughfare	
cascajo	dunul	grimstone	layoint	mysten	ruhe	tipper	
chappell	dwyer	gunbarrel	legault	nakai	ryark	tipperary	
chaseville	eachuston	hiwan	lincoln	nesda	san isabel	tivoli	
clark fork	ecklund	inavale	littlebear	newcomb	sawcreek	tomah	

DELAWARE

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

downer	fort mott	hammorton	hurlock	klej	plummer	rumford
evesboro	galestown	hooksan	kenansville	osier	pocomoke	

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL TYPE RESTRICTION TABLE (continued)

GEORGIA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. **At planting application only:** If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to **cotton, peanuts or soybeans** as an At-plant application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **700 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. **Side Dress Application Only:** If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to **cotton** as a Side Dress application only and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **700 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. **Split Application:** If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to **cotton** as an At-plant application and a Side Dress application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **1000 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. **At-planting and/or Post emergence application to peanuts:** If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to peanut as an At-plant application and/or a Post emergence (Pegging) application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **1100 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

americus	cainhoy	echaw	gorgas	kureb	meadowbrook	plummer	scranton
bigbee	centenary	eustis	hurricane	lakeland	molena	pottsburg	valdosta
blanton	chiple	flomaton	johns	leon	osier	ridgeland	wateree
boulogne	chipola	foxworth	kershaw	lynn haven	ousley	ridgewood	
buncombe	duckston	fripp	kingsferry	mandarin	penney	sapelo	

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL TYPE RESTRICTION TABLE (continued)

ILLINOIS

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

algansee	brems	coloma	granby	lanier	plainbo	seafield	watseka
alvin	burkhardt	dickinson	hodge	lorenzo	plainfield	sparta	zumbro
beavercreek	carmi	disco	homer	matherton	princeton	spinks	
billett	carr	el dara	hononegah	minneiska	robby	stonelick	
bloomfield	casco	eleva	hoopeston	morocco	rodman	thetford	
boone	chelsea	flagler	lamont	oakville	saffell		
boyer	chute	grable	landes	oshtemo	sarpy		

INDIANA

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

abskota	branch	coloma	hanna	maumee	ouiatenon	seafield	thetford
algansee	brems	conotton	homer	morocco	plainfield	shipshe	tyner
alvin	bronson	conrad	hononegah	moundhaven	princeton	sisson	wasepi
billett	bruno	desker	hoopeston	nesius	prochaska	sparta	watseka
bloomfield	carmi	dickinson	junius	newton	riverdale	spinks	zaborosky
bobtown	casco	elston	kosciusko	oakville	robby	stockland	zadog
boyer	chatterton	granby	landes	ormas	rodman	stonelick	
brady	chelsea	gravelton	leoni	oshtemo	saugatuck	tedrow	

IOWA

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

backbone	buckney	chute	estherville	granby	lamont	oesterle	sparta
billett	burkhardt	dickinson	finchford	hawick	landes	perks	toolesboro
boone	carr	dickman	flagler	hoopeston	lilah	salida	watseka
brady	chelsea	elrick	fruitfield	klum	montieth	sarpy	zenor

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL TYPE RESTRICTION TABLE (continued)

KANSAS

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

aline	dix	elsmere	gracemore	kanza	lincoln	schamber	tivoli
bankard	dwyer	goltry	haxtun	las animas	pratt	simeon	valent
dillwyn	els	goodnight	inavale	likes	sarpy	thurman	valentine

KENTUCKY

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

bruno	flomaton	iuka	lakin	molena	potomac	saffell	smithdale
crevasse							

LOUISIANA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

alaga	bienville	cheniere	eustis	hackberry	kenney	lotus	palm beach
bassfield	bigbee	crevasse	felicity	haggerty	lakeland	nugent	peveto
betis	bruno	darden	flo	hainesville	latonia	osier	

MARYLAND

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

aura	dragston	fort mott	hooksan	lakeland	newhan	remlik
carteret	duckston	fripp	hurlock	leetonia	osier	runclint
colts neck	eyesboro	galestown	ingleside	leon	plummer	
downer	fisherman	hammorton	klej	molena	pocomoke	

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL TYPE RESTRICTION TABLE (continued)

MICHIGAN

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

abbaye	brems	eleva	ishpeming	maumee	paavola	sayner	vilas
abscota	bronson	elston	kalkaska	mecosta	padus	seafield	wainola
adams	channing	ensign	karlin	menahga	pelkie	shawano	waiska
alcona	chatham	esau	kawbawgam	millersburg	pemene	shelldrake	wallace
algansee	chelsea	evart	keweenaw	monico	pence	sisson	wasepi
allouez	chesaning	fabius	kingsville	montcalm	pequaming	sparta	waseka
alpena	coloma	finch	kiva	morocco	perrin	spinks	winterfield
alvin	covert	freesoil	klacking	nadeau	peshekee	springlake	zeba
arcadian	croswell	gilchrist	koontz	newton	pipestone	st. ignace	zimmerman
arkport	cunard	gladwin	kosciusko	nottawa	plainfield	steuben	
au gres	deer park	granby	landes	oakville	rapson	sturgeon	
banat	deerton	grattan	leelanau	oconto	richter	summersville	
battlefield	desker	graycalm	leoni	ocqueoc	ridgeville	sundog	
bixler	dickinson	grayling	liminga	omega	riverdale	tedrow	
blue lake	dixboro	guardlake	lode	onota	rodman	tekenink	
bohemian	duel	halfaday	mahtomedi	ormas	rousseau	thetford	
boyer	dunbridge	hodenpyl	mancelona	oshtemo	rubicon	trimountain	
brady	east lake	ingalls	manitowish	otisco	sarona	tyre	
branch	eastport	ionia	matherton	ottokee	saugatuck	velvet	

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL TYPE RESTRICTION TABLE (continued)

MINNESOTA

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

abskota	burkhardt	deer park	granby	lasa	northfield	rosy	venlo
alban	burnsville	dickinson	graycalm	leola	nymore	salida	verndale
alganssee	carmi	dickman	hamar	lilah	oesterle	sanburn	wamduska
alvin	caryville	egeland	hawick	lino	omega	sartell	watseka
andrusia	chelsea	eleva	hecla	lohnes	osakis	serden	waupaca
anoka	chetek	emmert	hiwood	maddock	pierz	shawano	wawina
arvilla	claire	evart	hoopeston	mahtomedi	plainbo	sioux	winterfield
beavercreek	cloquet	faunce	insula	marquette	plainfield	soderville	zenor
becker	conic	finchford	kanaranzi	meehan	poppleton	sparta	zimmerman
bellchester	copaston	flagler	karlstad	menaha	quetico	spinks	zipfel
bergkeller	cormant	fossium	kost	mesaba	redby	sugarbush	zumbro
billett	cowhorn	friendship	lamont	minneiska	renshaw	torning	
boone	cromwell	gotham	langola	nemadji	rosholt	ulen	

MISSISSIPPI

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

alaga	beulah	bruno	eustis	lakeland	nugent	pactolus	saffell
bassfield	bigbee	crevasse	flomaton	latonia	osier	plummer	st. lucie

MISSOURI

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

alvin	canalou	crevasse	finchford	malden	sarpy	sparta
beulah	carr	diehlstadt	hodge	plainfield	scotco	wideman
bruno	clana	eustis	landes	sandbur	shelldrake	

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL TYPE RESTRICTION TABLE (continued)

MONTANA

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

alberton	biglake	colburn	glenberg	lisk	oceanet	ryorp	tamely	upsata
ambrant	blackhall	comad	glendive	lolopeak	oraid	sacheen	tassel	usine
apeldorn	blanchard	como	hanly	lone rock	ovando	sachett	terry	utica
ashbon	bonner	cowood	haskill	lowercreek	pend orielle	sawcreek	tinsley	valentine
ashuelot	brandenburg	cozberg	haverdad	macfarlane	poin	scravo	tipperary	vebar
assinniboine	branham	dast	haverly	manning	redcreek	selle	tomichi	victor
bangston	busby	dayschool	havrelon	mccaffery	rencot	selway	travessilla	virgelle
bankard	cabba	dominic	hiwan	mcilwaine	rentsac	seroco	travson	wabek
banks	calicott	dwyer	jugson	mires	rhame	sheege	treble	wilsonville
baxendale	castner	elkner	kalsted	mirror	ridge	shingle	trembles	woodgulch
bearmouth	cheadle	elmira	kirby	moiese	riedel	shook	trey	woodside
beaverton	cheyenne	ervide	krause	nelson	riverside	shurley	tulloch	worland
beenom	chinook	flasher	laporte	nemote	rivra	splitro	turnercrest	yellowbay
beisigl	clark fork	fleak	larim	nesda	rochester	stecum	tusler	yetull
belain	cohagen	glaciercreek	lihen	nippt	ryell	tally	twilight	zeona

NEBRASKA

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

alda	boone	craft	dwyer	hedville	manter	orwet	simeon	valentine
almeria	brunswick	dailey	eckley	hersh	marlake	phiferson	talmo	whitelake
anselmo	busher	darr	els	hoffland	mckelvie	pivot	tassel	wildhorse
bankard	calamus	dickinson	elsmere	holt	meadin	platte	thurman	
barney	canyon	dix	fonner	inavale	munjor	ronson	trelona	
bigwinder	carr	doger	gannett	ipage	o'neill	sarpy	treon	
blendon	cass	draknab	glenberg	laird	ord	schamber	tryon	
boel	chappell	duda	gothenburg	las animas	orpha	selia	tuthill	
bolent	cheyenne	dunday	haverson	lute	ortello	shingle	valent	

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL TYPE RESTRICTION TABLE (continued)

NORTH CAROLINA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

alaga	butters	conetoe	fripp	lakeland	nansemond	pactolus	rumford	valhalla
alpin	cainhoy	corolla	galestown	leon	newhan	plummer	seabrook	wake
baymeade	candor	dragston	immokalee	lillington	norfolk	pocomoke	st. lucie	wakulla
biltmore	carteret	duckston	johns	louisburg	ona	potomac	suncook	wando
blanton	centenary	echaw	kenansville	lynn haven	osier	pottsburg	tarboro	wateree
bojac	charleston	eustis	klej	mandarin	ostin	remlik	toisnot	
buncombe	chiple	foreston	kureb	molena	ousley	rimini	tomahawk	

NORTH DAKOTA

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

appam	blanchard	cormant	fossum	karlsruhe	minnewaukan	seroco	twilight	yecross
arvilla	brandenburg	dast	glendive	ladner	osakis	sioux	ulen	yetull
aylmer	brien	desart	hamar	lemert	poppleton	stirum	vebar	zeona
banks	cabba	egeland	hanly	lihen	renshaw	telfer	venlo	
bantry	chinook	ekalaka	haskill	lohnes	rhame	tinsley	verendrye	
beisigl	claire	falsen	havrelon	maddock	ruso	trembles	virgelle	
binford	coe	flasher	hecla	manning	schaller	trey	wabek	
blackhall	cohagen	fleak	inkster	metigoshe	serden	tuster	wamduska	

OHIO

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

abscota	bixler	coloma	dunbridge	homer	landes	otisville	princeton	spinks	tyner
algansee	boyer	colonie	elnora	hoopeston	lanier	ottokee	rodman	stafford	watertown
alvin	brady	conotton	galen	ionia	maumee	perrin	schaffenaker	steinsburg	
arkport	bronson	dekalb	granby	kingsville	oakville	pinegrove	sisson	stonelick	
barkcamp	casco	dixboro	hazleton	lakin	oshtemo	plainfield	sparta	tedrow	

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL TYPE RESTRICTION TABLE (continued)

SOUTH CAROLINA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. **At planting application only:** If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to **cotton, peanuts or soybeans** as an At-plant application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **700 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. **Side Dress Application Only:** If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to **cotton** as a Side Dress application only and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **700 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface. **Split Application:** If TEMIK brand 15G Lock 'n Load Aldicarb Pesticide is applied to **cotton** as an At-plant application and a Side Dress application and a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **1000 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

alaga	buffers	corolla	foreston	lakeland	nansemond	pocomoke	tarboro
alpin	cainhoy	crevasse	foxworth	leon	newhan	ridgeland	tomahawk
baratari	candor	dragston	fripp	louisburg	olanta	rimini	wakulla
baymeade	carteret	duckston	johns	lynn haven	osier	rosedhu	wando
blanton	centenary	echaw	kenansville	molena	ousley	scranton	waterree
brogdon	charleston	elloree	kershaw	mouzon	pactolus	seabrook	witherbee
buncombe	chipley	eustis	kureb	murad	plummer	seewee	

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL TYPE RESTRICTION TABLE (continued)

SOUTH DAKOTA

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

almeria	butche	duda	gannett	las animas	o'neill	shingle	tusler
alwilda	cabba	dunday	glenberg	lihen	orton	simeon	tuthill
anselmo	canyon	dwyer	glendive	lute	orwet	sioux	twilight
arvilla	cass	eckley	hamar	maddock	platte	stirum	ulen
assinniboine	chappell	egeland	hanly	manning	renshaw	storla	valent
aylmer	chinook	ekalaka	hecla	manter	reva	talmo	valentine
bankard	cohagen	els	henkin	marlake	rhame	tassel	vebar
banks	claire	elsmere	holt	mawer	rockoa	telfer	venlo
bantry	craft	evridge	hopdraw	mccaffery	ronson	thurman	wabek
beisigl	dailey	fedora	inavale	meadin	sarpy	travessilla	whitelake
blackhall	dix	flasher	ipage	minnewaukan	schamber	trembles	yecross
blendon	doger	fleak	kirby	munjor	serden	trey	zeona
boel	dogiecreek	fossum	ladner	murdo	seroco	tryon	

TENNESSEE

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

barkcamp	bigbee	bruno	buncombe	crevasse	eustis	nugent	potomac
----------	--------	-------	----------	----------	--------	--------	---------

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL TYPE RESTRICTION TABLE (continued)

VIRGINIA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

alaga	camocca	dragston	johns	lakin	nansemond	pocomoke	tarboro
assateague	carteret	drall	kenansville	leetonia	newhan	remlik	wakulla
aura	catpoint	duckston	klej	leon	osier	rumford	wando
biltmore	chipley	fisherman	kureb	magotha	ousley	schaffenaker	wateree
bojac	conetoe	fripp	lakehurst	millrock	pactolus	seabrook	
buncombe	corolla	galestown	lakeland	molena	plummer	spessard	

WEST VIRGINIA

If a vulnerable soil is present and the water table is less than 25 feet below ground surface, do not apply within **300 feet** of a drinking water well unless it is known or reasonably believed based upon authoritative sources that such wells are either cased to 100 feet below ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

drall	lakin	landes	leetonia	potomac	schaffenaker
-------	-------	--------	----------	---------	--------------

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"

SOIL TYPE RESTRICTION TABLE (continued)

WYOMING

If a vulnerable soil is present and the water table is less than 25 feet, do not apply within **500 feet** of a drinking water well unless it is known or reasonably believed based upon an authoritative source that such wells are either cased to 100 feet below the ground level or a minimum of 30 feet below the water table. If it is not known whether the water table is greater than 25 feet below ground surface, assume that the water table is less than 25 feet below ground surface.

ashbon	calicott	dunday	haverly	millburne	ratake	stecum	twilight
adllis	cambarge	dunul	havermom	milok	redbank	sudworth	valent
alcova	canwall	dwyer	hazton	mishak	redcreek	sunup	valentine
alflack	carbol	eckley	hechtman	mosroc	redfeather	taluce	vible
anasazi	cass	edlin	henrieville	motoqua	rekop	tassel	vonason
anselmo	castner	elk mountain	herbman	mudray	relob	tasselman	walcott
apeldorn	cathedral	ellicott	hiland	nathale	rencot	teagulf	walden
assinniboine	cestnik	emblem	hiwan	nelson	reno	teewinot	wall
bankard	chappell	enos	hoot	nesda	rentsac	tellman	wardboro
banks	cheadle	ethelman	huguston	newfork	rivra	tenorio	wendover
barcus	chinook	eyre	irigul	nizina	rogert	terada	wetmore
barnum	clarkelen	farisita	julesburg	norrison	rohonda	terro	whaley
baroid	comad	farson	kandaly	norte	roxal	terry	wiggleton
barrett	conpeak	feltner	koonich	oceanet	ryark	tetonville	willwood
bearmouth	cordes	fleak	labou	onason	ryorip	theedle	wilsonville
beaverton	corlett	folavar	lamarsh	orpha	sachett	thermopolis	wint
bigwin	cotha	frontier	lambman	otterson	sawcreek	tieside	worf
bigwinder	cotopaxi	gas creek	laney	ouray	schamber	tine	worfka
billycreek	cowestglen	glenberg	laporte	pahlow	schooner	tipper	worfman
blackhall	cowood	glendive	lariat	peetz	scravo	tipperary	worfstone
bobtail	coyet	glenton	larim	pendergass	seedskaadee	tolman	worland
bodorumpe	cragosen	grenoble	las animas	pepton	sharlard	travessilla	yetull
bosler	crestman	grieves	lazear	pescar	sheege	travson	youjay
bottle	crownest	grimm	littlebear	phiferson	shingle	trelona	zeomont
boyle	cryluha	grimstone	littsan	pilotpeak	shoshone	trembles	zeona
branham	curabith	gunbarrel	lone rock	pineguest	shurley	treoff	
brece	dailey	handran	macfarlane	poin	sobson	treon	
brownsto	dast	hanks	manter	preatorson	southace	trook	
bruja	derrick	hanly	mathers	pugsley	southfork	troutville	
busher	devore	hapjack	mccaffery	quealman	space city	tullock	
butche	dix	haterton	meadowlake	quealy	splitro	turnback	
bymie	draknab	haverdad	means	randman	spool	turnercrest	

US79862768D TEMIK 15 G LOCK 'N LOAD ETL

5.75" x 3.875"