



**FOR CONTROL OF CERTAIN BROADLEAF WEEDS IN SMALL GRAINS (WHEAT, BARLEY, OATS AND RYE), CONSERVATION RESERVE PROGRAM (CRP) AREAS, GRASSES GROWN FOR SEED OR SOD PRODUCTION AND FLAX**

**ACTIVE INGREDIENTS:**

Octanoic acid ester of bromoxynil (3,5-dibromo-4-Hydroxybenzonitrile)*	31.7%
2-Ethylhexyl ester of 2-methyl-chlorophenoxyacetic acid**	34.0%
<b>OTHER INGREDIENTS***</b>	<b>34.3%</b>
<b>TOTAL:</b>	<b>100.0%</b>

\* Bromoxynil octanoate equivalent to 21.8% of bromoxynil or not less than 2.0 pounds of bromoxynil per gallon.

\*\* Equivalent to 21.8% 2-methyl-chlorophenoxyacetic acid or not less than 2.0 pounds MCPA acid per gallon.

\*\*\*Contains petroleum distillates

EPA Reg. No. 2935-552

☐ EPA Est. No. 228-IL-1

☐ EPA Est. No. 42750-MO-1

**KEEP OUT OF REACH OF CHILDREN  
WARNING - AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>NOTE TO PHYSICIAN:</b> Contains petroleum distillate, vomiting may cause aspiration pneumonia.	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**Warning:** May be fatal if swallowed. Harmful if absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist.

**SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL  
PRECAUTIONARY STATEMENTS.**

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical resistant to this product are made of barrier laminate, butyl rubber, nitrile rubber or viton. If you want more options, follow the instructions for category F on an EPA chemical-resistant category selection chart.

**Mixers, loaders, applicators, flaggers, and other handlers must wear:** Coveralls over a long-sleeved shirt and long pants, chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, or viton gloves, a chemical-resistant apron when cleaning equipment, protective eyewear, chemical-resistant headgear for overhead exposure, and chemical-resistant footwear plus socks. In addition to the above, mixers, loaders, and cleaners of equipment must also wear chemical-resistant gloves and a chemical-resistant apron. Mixers and loaders supporting aerial application must also wear a NIOSH approved particulate filtering respirator equipped with N, R, or P class filter media. The respirator should have a NIOSH approval number prefix TC-84A. It is recommended the respirator wearer be fit tested, and trained in the use, maintenance, and limitations of the respirator. See "Engineering Controls" for additional requirements.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROLS**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard(WPS) for agricultural pesticides [40CFR 70.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. **IMPORTANT:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(4-6)].

If you will handle a total of 60 gallons or more of this product per day, you must use a mechanical transfer system for all mixing and loading operations. If this product is packaged in a 30 gallon drum, you must use a mechanical transfer system which terminates in a drip-free hard coupling which may be used only with a spray or mix tank which has been fitted with a compatible coupling. If you do not presently own or have access to a mechanical transfer system with this type of coupling, contact your dealer for information on how to obtain such a system or to modify your present system.

When using a mechanical transfer system, do not remove or disconnect the pump or probe from the container until the container has been emptied and rinsed. The pump or probe system must be used to rinse the empty container and to transfer the rinsate directly to the mixing or spray tank.

Application from a tractor with a completely enclosed cab or aerial application is required whenever this product is applied to 360 or more acres in a day. The closed systems and enclosed cabs must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)]. The handler PPE requirements may be reduced or modified as specified in the WPS.

To reduce exposure to residues, wash the spray rig, tractor, and all other equipment used to handle or apply this product with water daily or before using the equipment for any other purpose.

APPLICATION BY CHEMIGATION must be done by fixed pipe, overhead sprinkler systems or hand-moved pipe. If hand-moved pipe is used for chemigation, the pipe must not be handled in any way until 24 hours after chemigation has been completed and residues have been flushed from the system. When applying by chemigation, no person may enter the application site unless in an enclosed vehicle.

DURING AERIAL APPLICATION, human flaggers are prohibited unless in enclosed vehicles. Aerial application is prohibited within 300 feet of residential areas (e.g., homes, schools, hospitals, shopping areas, etc.)

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish, aquatic invertebrates and aquatic plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. This pesticide has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

### PHYSICAL AND CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

### IMPORTANT

Vendetta® Herbicide contains low volatile 2-ethylhexyl ester of MCPA. At high air or ground surface temperatures, vapors from Vendetta Herbicide may cause injury to susceptible plants. This fact should be considered when applying Vendetta Herbicide.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Use of this product in certain portions of California, Oregon and Washington is subject to the January 22, 2004 Order for injunctive relief with Washington Toxics Coalition, et. al. v. EPA C01-0132C, (W.D. WA). For further information, please refer to <http://www.epa.gov/espp/litstatus/wtc/index.htm>

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI). For all crops except turf, the REI is 24 hours. The REI for harvesting sod farm turf is 26 days. The REI for other turf activities is 24 hours. For uses on turf grown for transplanting (e.g. on sod farms), notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is chemical-resistant gloves made of any waterproof material, coveralls, shoes plus socks and protective eyewear.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to the use of this product on non-residential turfgrass areas that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

### USE INFORMATION

Vendetta Herbicide is formulated as an emulsifiable concentrate containing the equivalent of 2 lbs. per gallon of bromoxynil and 2 pounds per gallon of 2-ethylhexyl ester of MCPA.

Vendetta Herbicide is a selective postemergence herbicide for control of important broadleaf weeds infesting small grains (wheat, barley, oats, rye), flax, conservation reserve program areas, and grass grown for seed or sod production. Optimum weed control is obtained when Vendetta Herbicide is applied to actively growing weed seedlings. Vendetta Herbicide is primarily a contact herbicide, therefore thorough coverage of the weed seedlings is essential for optimum control.

Vendetta Herbicide has little residual activity. Therefore subsequent flushes of weeds will not be controlled by the initial treatment. Generally crops that form a good canopy will help shade subsequent weed flushes. However, certain crops or short-straw varieties, for example Yecora Rojo wheat, may not develop the crop canopy fast enough to shade the subsequent flushes of weeds.

Occasional transitory leaf burn may occur. The temporary leaf burn is similar to that seen with liquid fertilizer. Because the activity of Vendetta Herbicide is mainly contact, recovery of the crop is generally rapid with no lasting effect. Frequency and amount of leaf burn may be greater when crops are stressed by abrasive winds, cool to cold evening temperatures or mechanical injury, such as that caused by hail, sleet or insect feeding. To reduce the potential for temporary leaf burn, applications should be made to dry foliage in the recommended spray volumes per acre when weather conditions are not extreme.

## MIXING, LOADING AND HANDLING INSTRUCTIONS

### 2.5 Gallon Containers

Special care must be taken in mixing and loading Vendetta Herbicide. Hands should be placed on the container in such a way as to avoid possible drip or splash.

### 30 Gallon and Bulk Containers

If you will handle a total of 60 gallons or more of Vendetta Herbicide per day, you must use a mechanical transfer system for all mixing and loading operations. If Vendetta Herbicide is packaged in a 30 gallon drum, you must use a mechanical transfer system which terminates in a drip-free hard coupling which may be used only with a spray or mix tank which has been fitted with a compatible coupling. If you do not presently own or have access to a mechanical transfer system with this type of coupling, contact your dealer for information on how to obtain such a system or to modify your present system.

When using a mechanical transfer system, do not remove or disconnect the pump or probe from the container until the container has been emptied and rinsed. The pump or probe system must be used to rinse the empty container and to transfer the rinsate directly to the mixing or spray tank.

**VENDETTA HERBICIDE ALONE:** Fill the spray tank 1/2 to 3/4 full with clean water. Begin agitation and add the recommended amount of Vendetta Herbicide. Add water to the spray tank to the desired level. Maintain sufficient agitation to ensure a uniform spray mixture during application.

**TANK MIXTURES:** Vendetta Herbicide can be applied in tank mixture with many other herbicides and insecticides registered for use on approved crops. Refer to the specific crop section for rate recommendations and other restrictions. To apply Vendetta Herbicide in mixture with another product, fill the spray tank 1/2 to 3/4 full with clean water and begin agitation. If tank mixing with wettable powder, soluble powder, flowable or dry flowable products, add the powder or flowable product first. After the other herbicide is thoroughly mixed with water, add the specified amount of Vendetta Herbicide and add water to the spray tank to the desired level. If tank mixing with other product types, add Vendetta Herbicide first before adding the other product. Always mix one product in water thoroughly before adding another product or compatibility problems may occur. Never mix two products together without first mixing in water.

Maintain sufficient agitation while mixing and during application to ensure a uniform spray mixture. If spray mixture is allowed to remain without agitation for short periods of time, be sure to agitate until uniformly mixed before application.

If tank mixing with products other than those listed within each crop section, a compatibility test is recommended to ensure satisfactory spray preparation. To test for compatibility, use a small container and mix a small amount (0.5 to 1 quart) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5 to 15 minutes after mixing. To ensure maximum crop safety and weed control, follow all cautions and limitations on this label and the labels of products used in the tank mixture with Vendetta Herbicide.

## SPRAYABLE LIQUID FERTILIZERS AND SPRAY ADDITIVES

Vendetta Herbicide can be applied in combination with sprayable liquid fertilizer or spray additives such as surfactants or crop oil concentrate. When tank mixing with liquid fertilizer always add the fertilizer to the spray tank first and agitate thoroughly before adding Vendetta Herbicide. Always predetermine the compatibility with liquid fertilizer by mixing small proportional quantities in advance. Agitation must be maintained during filling and application operations to ensure that Vendetta Herbicide is evenly mixed with the fertilizer. Leaf burn may occur when Vendetta Herbicide is applied with liquid fertilizer, but new leaves are not adversely affected.

**IMPORTANT:** Fertilizers and spray additives can increase foliage leaf burn when applied with Vendetta Herbicide. Do not apply fertilizers or spray additives with Vendetta Herbicide if leaf burn is a major concern due to environmental conditions, crop or variety sensitivity to Vendetta Herbicide.

## APPLICATION PROCEDURES

Vendetta Herbicide can be applied to registered use areas by ground, aerial and sprinkler irrigation equipment.

### GROUND APPLICATION

Use a standard herbicide boom sprayer that provides uniform and accurate application. Sprayer should be equipped with screens no finer than 50 mesh in the nozzle tips and in-line strainers.

Select a spray volume and delivery system that will ensure thorough and uniform spray coverage. For optimum spray distribution and thorough coverage use of flat fan nozzles (maximum tip size 8008) with a spray pressure of 40-60 psi are recommended.

Other nozzle types and lower spray pressures that produce coarse spray droplets may not provide adequate coverage of the weeds to ensure optimum control. Raindrop® nozzles and flood nozzles are not recommended as weed control with Vendetta Herbicide may be reduced.

In general, a spray volume of 10 to 20 gallons per acre (GPA) is recommended for optimum spray coverage. A minimum of 5 GPA with a minimum spray pressure of 50 psi and a maximum ground speed of 10 mph may be used with higher speed, low volume ground application if ground terrain, crop and weed density allow effective spray distribution. When using higher speed equipment, a maximum ground speed of 10 mph is suggested if field conditions cause excessive boom movement during application which results in poor spray coverage. Ground applications made when dry, dusty field conditions exist may provide reduced weed control in wheel track areas. Applications using less than 10 gallons per acre may result in reduced weed control.

When weed infestations are heavy, use of higher spray volumes and spray pressure will be helpful in obtaining uniform weed coverage. When corn or grain sorghum are large enough to interfere with the spray pattern, drop nozzles should be used to obtain uniform weed coverage. If you are unsure of the infestation level or size of crop, consult your local extension service.

Do not apply when winds are gusty or when other conditions favor poor spray coverage and/or off target spray movement.

Do not apply with nozzle height greater than 4 feet above crop canopy.

### AERIAL APPLICATION

Use orifice discs, cores and nozzle types and arrangements that will provide for optimum spray distribution and maximum coverage. In general a minimum spray volume of 5 GPA and a maximum pressure of 40 psi are recommended. A minimum spray volume of 3 gallons per acre may be used if crop canopy and weed density allow adequate spray coverage. Aerial applications using less than 5 gallons of spray volume per acre may result in reduced weed control.

Do not apply during inversion conditions, when winds are gusty or when other conditions favor poor spray coverage and/or off-target spray movement. Off-target spray movement can be minimized by increasing the spray volume per acre and not applying when winds exceed 10 mph.

Nozzles must always point backward parallel with the airstream and never point downwards more than 45 degrees.

### SPRINKLER IRRIGATION APPLICATION

Vendetta Herbicide can be applied through sprinkler irrigation systems to small grains and grasses grown for seed or sod production.

Apply Vendetta Herbicide through sprinkler systems including center pivot, lateral move, side (wheel) roll, solid set or hand move irrigation systems only. If hand moved pipe is used for chemigation, the pipe must not be handled in any way until 24 hours after chemigation has been completed and residues have been flushed from the system. When applying by chemigation, no person may enter the application site unless in an enclosed vehicle. Do not apply Vendetta Herbicide through any other type of irrigation system.



## SPECIFIC REQUIREMENTS FOR APPLICATION THROUGH AUTOMATED SPRINKLER IRRIGATION SYSTEM

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Agitation is recommended in the pesticide supply tank when applying Vendetta Herbicide.
9. Vendetta Herbicide should be applied continuously for the duration of the water application with center pivot and continuous lateral move systems. Application of Vendetta Herbicide should be made during the last 30-45 minutes of the irrigation set with other overhead sprinkler systems.
10. For best performance, set the sprinkler system to deliver approximately 0.5 inch or less of water per acre.
11. Remove scale, pesticide residues and other foreign matter from the supply tank and entire injector system. Flush with clean water.
12. If Vendetta Herbicide is diluted in the supply tank, fill the tank with half of the water amount desired, add Vendetta Herbicide and then add remaining water amount with agitation. Always dilute with at least 4 parts water to 1 part Vendetta Herbicide.
13. Start the sprinklers and then inject Vendetta Herbicide into the irrigation line. Vendetta Herbicide should be injected with a positive displacement pump into the main line at least 8 feet ahead of a right angle turn to insure adequate mixing. Refer to Vendetta Herbicide label for detailed information on application rates and timings.

## CHEMIGATION USER PRECAUTIONS

Application of more than 0.5 inch/acre of irrigation water may result in decreased product performance on certain soils.

Do not apply when conditions favor drift, when system connections or fittings leak, or when nozzles do not provide uniform distribution.

Allow sufficient time for pesticide to be flushed through all the lines and nozzles before turning off irrigation water.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

Do not connect an irrigation system used for pesticide application to a public water system.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operations, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

## SPRAY DRIFT MANAGEMENT

**SENSITIVE AREAS:** The pesticide must only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitats for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when wind speed is 2-10 mph at the application site.

The following drift management requirements must be followed to avoid off target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulation.

### FOR AERIAL APPLICATION:

- The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
- Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.
- Apply only when wind speed is 2-10 mph at the application site.

### FOR GROUND BOOM APPLICATION:

- Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Where states have more stringent regulations, they must be observed. The applicator should be familiar with and take into account the information covered in the [Aerial Drift Reduction Advisory Information](#).

**INFORMATION ON DROPLET SIZE:** (This section is advisory in nature and does not supersede the mandatory label requirements).

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions below).

**CONTROLLING DROPLET SIZE:** (This section is advisory in nature and does not supersede the mandatory label requirements.)

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the air stream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

**BOOM LENGTH:** (This section is advisory in nature and does not supersede the mandatory label requirements.)

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

**APPLICATION HEIGHT:** (This section is advisory in nature and does not supersede the mandatory label requirements.)

Applications must not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

**SWATH ADJUSTMENT:** (This section is advisory in nature and does not supersede the mandatory label requirements.)

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.)

**WIND:** (This section is advisory in nature and does not supersede the mandatory label requirements.)

Drift potential is lowest between wind speeds of 2 - 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application must be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect spray drift.

**TEMPERATURE AND HUMIDITY:** (This section is advisory in nature and does not supersede the mandatory label requirements.)

When making application in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**TEMPERATURE INVERSIONS:** (This section is advisory in nature and does not supersede the mandatory label requirements.)

Applications must not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

## WEED LIST

Postemergence application of Vendetta Herbicide will control the following weeds when sprayed in the seedling stage. Maximum weed stage of growth are listed under the Vendetta Herbicide Directions tables.

### MOST SUSCEPTIBLE BROADLEAF WEED SPECIES

Annual sowthistle	( <i>Sonchus oleraceus</i> )
Black mustard	( <i>Brassica nigra</i> )
Black nightshade	( <i>Solanum nigrum</i> )
Common cocklebur	( <i>Xanthium strumarium</i> )
Common lambsquarters	( <i>Chenopodium album</i> )
Common tarweed	( <i>Hemizonia congesta</i> )
Cow cockle	( <i>Saponaria vaccaria</i> )
Cutleaf nightshade	( <i>Solanum triflorum</i> )
Eastern black nightshade	( <i>Solanum ptycanthum</i> )
Coast fiddleneck	( <i>Amsinckia intermedia</i> )
Field pennycress	( <i>Thlaspi arvense</i> )
Green smartweed	( <i>Polygonum scabrum</i> )
Hairy nightshade	( <i>Solanum sarachoides</i> )
Horned Poppy	( <i>Glaucium corniculatum</i> )
Jimsonweed	( <i>Datura stramonium</i> )
Ladysthumb	( <i>Polygonum persicaria</i> )
Lanceleaf sage	( <i>Salvia reflexa</i> )
London rocket	( <i>Sisymbrium irio</i> )
Marshelder	( <i>Iva xanthifolia</i> )

### MOST SUSCEPTIBLE BROADLEAF WEED SPECIES(continued)

Pennsylvania smartweed	( <i>Polygonum strumarium</i> )
Pepperweed spp.	( <i>Lepidium app.</i> )
Redroot pigweed	( <i>Amaranthus retroflexus</i> )
Russian thistle	( <i>Salsola kali</i> )
Shepherdspurse	( <i>Capsella bursa-pastoris</i> )
Silverleaf nightshade	( <i>Solanum elaeagnifolium</i> )
Smooth pigweed	( <i>Amaranthus hybridus</i> )
Spiny pigweed	( <i>Amaranthus spinosus</i> )
Sunflower <sup>1</sup>	( <i>Helianthus annuus</i> )
Tall Waterhemp	( <i>Amaranthus tuberculatus</i> )
Tartary buckwheat	( <i>Fagopyrum tataricum</i> )
Tumble mustard	( <i>Sisymbrium altissimum</i> )
Wild buckwheat	( <i>Polygonum convolvulus</i> )
Wild mustard	( <i>Sinapis arvensis</i> )
Yellow rocket	( <i>Barbarea vulgaris</i> )

<sup>1</sup> For control of sunflower, delay application until first sunflower seedlings emerging are 4 inches in height.

### SUSCEPTIBLE BROADLEAF WEED SPECIES

Blue (purple) mustard	( <i>Chlorispora tenella</i> )
Common groundsel	( <i>Senecio vulgaris</i> )
Common ragweed	( <i>Ambrosia artemisiifolia</i> )
Corn chamomile	( <i>Anthemis arvensis</i> )
Corn gromwel	( <i>Lithospermum arvense</i> )
Fumitory	( <i>Fumaria officinalis</i> )
Giant ragweed	( <i>Ambrosia trifida</i> )
Hemp sesbania	( <i>Sesbania exaltata</i> )
Henbit	( <i>Lamium amplexicaule</i> )
Ivyleaf morningglory	( <i>Ipomoea hederacea</i> )
Knawel	( <i>Scleranthus annuus</i> )
Kochia	( <i>Kochia scoparia</i> )
Mayweed	( <i>Anthemis cotula</i> )
Prostrate knotweed	( <i>Polygonum aviculare</i> )
Puncture vine	( <i>Tribulus terrestris</i> )
Tall morningglory	( <i>Ipomoea purpurea</i> )
Tansy mustard	( <i>Descurainia pinnata</i> )
Tarweed	( <i>Hemizonia spp.</i> )
Velvetleaf	( <i>Abutilon theophrasti</i> )
Wild radish	( <i>Raphanus raphanistrum</i> )

Weeds germinating after spraying will not be controlled.

### WEED SUPPRESSION

Canada Thistle	( <i>Cirsium arvense</i> )
----------------	----------------------------

Vendetta Herbicide applied at 1-1/2 pints per acre provides burn down of top growth. Regrowth may occur. Make applications when Canada thistle is 8 inches tall to the bud stage.

**WHEAT, BARLEY, OATS AND RYE  
VENDETTA HERBICIDE DIRECTIONS**

		APPLICATION TIMING AND SPECIFIC COMMENTS	
PRODUCT	RATE	CROP	WEEDS
VENDETTA HERBICIDE	1 pint/A	Fall seeded wheat, barley, oats and rye throughout the United States and spring seeded wheat, barley, oats and rye in Idaho, Oregon, Washington, Colorado, Wyoming and Montana.  Apply to wheat, barley, oats and rye from the 3-leaf stage but before the crop reaches the boot stage.	MOST SUSCEPTIBLE BROADLEAF WEEDS: Apply to weeds up to the 8-leaf stage or 4 inches in height, whichever comes first. If weed forms rosette, apply before weeds exceed 2 inches in diameter.
	1-1/2 to 2 pints/A		SUSCEPTIBLE BROADLEAF WEEDS: Apply to weeds up to the 4-leaf stage or 2 inches in height, whichever comes first. If weed forms rosette, apply before weeds exceed 1 inch in diameter.
	2 pints/A		Apply to henbit, knawel and mayweed up to the 4-leaf stage or 2 inches in height, whichever comes first. Apply to kochia and tansy mustard for improved control when these weeds exceed the recommended stage of growth or are growing under cool, dry conditions.
	1-1/2 pints/A	Spring seeded wheat and barley except Idaho, Oregon, Washington, Colorado, Montana, and Wyoming. Apply to wheat, barley, oats and rye from the 3-leaf stage but before the crop reaches the boot stage.	MOST SUSCEPTIBLE AND SUSCEPTIBLE BROADLEAF WEEDS: Apply to weeds that do not exceed the 8-leaf stage or 4 inches in height, whichever comes first. If weed forms rosette, apply before weeds exceed 2 inches in diameter. Apply to kochia up to 2 inches in height.
	1-1/2 to 2 pints/A	Spring seeded wheat and barley except Idaho, Oregon, Washington, Colorado, Montana, and Wyoming. Apply to wheat, barley, oats and rye from the 3-leaf stage but before the crop reaches the boot stage.	Apply to kochia that is 2-4 inches height.
	Chemigation Only 2 pints/A	Apply to wheat, barley, oats and rye from the 3-leaf stage but before the boot stage. Apply through automated sprinkler irrigation systems with mechanical transfer loading system only. See MIXING LOADING AND HANDLING INSTRUCTIONS section for complete details.	Apply to MOST SUSCEPTIBLE and SUSCEPTIBLE broadleaf weeds up to the 4-leaf stage, 2 inches in height or 1 inch in diameter, whichever comes first.
	Post-harvest 3/4 to 2 pints/A	Make applications following harvest of wheat, barley, oats and rye in the states of North Dakota, South Dakota, Minnesota, and Montana. Do not plant any rotational crop until the following use season.	Apply 3/4 to 1 pint/A to MOST SUSCEPTIBLE BROADLEAF WEEDS up to the 8-leaf stage or 4 inches in height, whichever comes first. Apply 1-1/2 to 2 pints/A to SUSCEPTIBLE BROADLEAF WEEDS up to the 4-leaf stage or 2 inches in height, whichever comes first. For control of both grasses and broadleaf weeds, tank mix Vendetta Herbicide with Roundup® or Roundup® + 2,4-D such as WEEDONE® or WEEDAR® brand herbicides.

**WHEAT, BARLEY, OATS AND RYE**  
**VENDETTA HERBICIDE TANK MIXTURE DIRECTIONS**

APPLICATION TIMING AND SPECIFIC COMMENTS			
PRODUCT	RATE	CROP	WEEDS
VENDETTA Herbicide + MCPA Ester	3/4 to 2 pints/A + 1/4 to 1/2 pint/A	Apply to spring seeded wheat, barley, oats and rye from tillering stage, but before boot stage.	For control of MOST SUSCEPTIBLE and SUSCEPTIBLE weeds and improved control of redroot pigweed and kochia. Apply to weeds up to the 8-leaf stage, 3 inches in height or 2 inches in diameter, whichever comes first. Apply to kochia and redroot pigweed up to 2 inches in height or diameter.
VENDETTA Herbicide + Glean® + nonionic surfactant	3/4 to 1-1/2 pints/A + 1/6 to 1/3 oz/A + 1 qt/100 gal. of water	Apply to wheat and barley from the 3-leaf stage but before the crop reaches the boot stage. Refer to Glean® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, tansy mustard and chickweed. Apply to weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.
VENDETTA Herbicide + Finesse® + nonionic surfactant	3/4 to 1-1/2 pints/A + 1/6 to 1/3 oz/A + 1 qt/100 gal. of water	Apply to wheat and barley from the 3-leaf stage but before the crop reaches the boot stage. Refer to Finesse® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, tansy mustard and chickweed. Apply to weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.
VENDETTA Herbicide + Ally® + nonionic surfactant	3/4 to 1-1/2 pints/A + 1/10 oz/A + 1 qt/100 gal. of water	Apply to wheat and barley from the 3-leaf stage but before the crop reaches the boot stage. Refer to Ally® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, tansy mustard and chickweed. Apply to weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.
VENDETTA Herbicide + Dicamba DMA Salt	3/4 to 1-1/2 pints/A + 1/8 to 1/4 pint/A	<b>FOR USE ON WHEAT ONLY. DO NOT TREAT BARLEY, OATS OR RYE.</b> Fall seeded wheat from the 3-leaf stage but before jointing. Spring seeded wheat from the 3 to 5 leaf stage of growth.	This tank mix improves control of broadleaves such as prostrate knotweed and kochia. Apply to weeds up to the 8 leaf stage, 3 inches in height or 2 inches in diameter, whichever comes first. Apply to kochia up to 2 inches in height or diameter.
VENDETTA Herbicide + Harmony® Extra + nonionic surfactant	3/4 to 1-1/2 pints/A + 3/10 to 1/2 oz/A + 1 qt/100 gal. of water	Winter wheat. Apply from the 3-leaf stage but before the 3rd node is detectable. Refer to the Harmony® Extra label for crop rotation and other restrictions.  Spring wheat and barley. Apply after the 3-leaf stage but before the 1st node is detectable. Refer to the Harmony® Extra label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, chickweed and redroot pigweed. Apply to weeds up to the 8-leaf stage, 4 inches in height or across, whichever comes first.
VENDETTA Herbicide + Amber® + nonionic surfactant	3/4 to 1-1/2 pints/A + 0.28 to 0.56 oz/A + 0.25 to 0.5% v/v	Apply to wheat and barley from the 3-leaf stage, but before the flag leaf is visible. Refer to the Amber® label for crop rotation and other restrictions.	This tank mix improves control of broadleaves such as henbit, tansy mustard, and pigweed. Apply to weeds up to the 4-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.
VENDETTA Herbicide + Express® + nonionic surfactant	3/4 to 1-1/2 pints/A + 1/6 to 1/3 oz/A + 1 qt/100 gal. of water	Wheat and barley. Apply from the 3-leaf stage but before the flag is visible. Refer to the Express® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, chickweed, redroot pigweed and suppression of Canada thistle. Apply to annual weeds up to the 8-leaf stage, 4 inches in height or across, whichever comes first and to Canada thistle 4 to 8 inches tall with 2 to 6 inches of new growth.



**WHEAT, BARLEY, OATS AND RYE**  
**VENDETTA HERBICIDE TANK MIXTURE DIRECTIONS (continued)**

APPLICATION TIMING AND SPECIFIC COMMENTS			
PRODUCT	RATE	CROP	WEEDS
VENDETTA Herbicide + Curtail® Curtail® M	3/4 to 1-1/2 pints/A + 2 pints/A	Apply to wheat and barley after the crop begins to tiller up to the 1st node detectable.	This tank mix improves control of kochia, wild buckwheat and suppression of Canada thistle. Apply to annual broadleaf weeds up to the 8 leaf stage, 4 inches in height or 2 inches in diameter and to Canada thistle in the rosette to prebud stage.
VENDETTA Herbicide + metribuzin (Sencor® or Lexone®)	1 pint/A + 1/8 to 3/16 lb. ai/A	Winter wheat in Idaho, Oregon and Washington. Apply in spring after growth has started and secondary roots with a minimum of 3 to 4 tillers have been established, but before the forming of joints in the stem. Avoid application when crop has experienced winter kill, frost damage, disease or drought.	This tank mix improves control of broadleaf weeds such as chickweed, filaree, henbit. Apply to weeds up to the 4 leaf stage, 2 inches in height or diameter, whichever comes first. A recognized authority should be consulted concerning the use of this mixture in your area.
VENDETTA Herbicide + Avenge®	1 to 2 pints/A + 2-1/2 to 4 pints/A	Winter wheat. Four leaf to tillering stage. Refer to Avenge® label for varietal and other restrictions.  Spring Wheat. Five to 6 leaf stage. Refer to Avenge® label for varietal and other restrictions.  Barley. Three to 7 leaf stage.	This tank mix will provide wild oat control in addition to broadleaves. Apply to wild oats in the 3-5 leaf stage and broadleaves that do not exceed the 4 leaf stage or rosettes of 1.5 inches in diameter. Avenge® use rates per acre are 2-1/2 pints (1-10 oats per sq. ft.), 3 pints (11-25 oats per sq. ft.) or 4 pints (more than 25 oats per sq. ft.).
VENDETTA Herbicide + Assert®	1 to 1-1/2 pints/A + 1 to 1-1/2 pints/A	Apply to wheat and barley from the 3-leaf stage but before boot stage. Refer to Assert® label for crop rotation and other restrictions.	This tank mix will provide wild oat control in addition to broadleaf weeds. Apply to wild oats at the 1-4 leaf stage and broadleaf weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first. Use Assert at 1-1/2 pints/A west of the Rocky Mountains or if wild oats have initiated tillering. For spray volumes in excess of 10 GPA, add 0.3 fluid oz of nonionic surfactant for each gallon in excess of 10 GPA.

**RESTRICTIONS and PRECAUTIONS: Wheat, Barley, Oats and Rye**

- Do not graze treated fields within 45 days after application.
- Do not apply when crops are under moisture stress.
- Do not apply when crop canopy covers the weeds as poor control will result.
- Reduced weed control may occur when weeds are stressed from lack of moisture or cold temperatures.
- Refer to labels of products used in tank mixture for additional restrictions and precautions.
- Do not apply more than 2 pints of Vendetta Herbicide (0.5 lb. MCPA acid equivalent) per acre in a single growing season.
- Do not plant rotational crops within 30 days following application of Vendetta Herbicide.
- Do not apply more than 0.75 lb. of MCPA acid equivalent per acre per year when applying alone or tank mixing with other products that contain MCPA.

**CONSERVATION RESERVE PROGRAM AREAS (CRP)**  
**VENDETTA HERBICIDE DIRECTIONS**

APPLICATION TIMING AND SPECIFIC COMMENTS			
PRODUCT	RATE	CROP	WEEDS
VENDETTA Herbicide	1 to 2 pints/A	Apply to grasses from the 3 leaf stage.	Apply 1 pint/A to MOST SUSCEPTIBLE and 1-1/2 to 2 pints/A to SUSCEPTIBLE broadleaf weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.

**RESTRICTIONS and PRECAUTIONS: CRP Areas**

- Do not allow livestock to graze in treated areas or feed treated grass to livestock.
- If legumes are included in CRP area planting, severe injury may occur to legumes treated with Vendetta Herbicide.
- Do not apply Vendetta Herbicide to CRP areas planted with alfalfa if temperatures are expected to exceed 80° F or severe crop injury may occur. If legumes other than alfalfa have been planted, severe crop injury may occur at any application temperature.
- Do not apply more than 2 pints/A of Vendetta Herbicide to CRP areas that are underseeded with alfalfa.



**GRASSES GROWN FOR SEED PRODUCTION OR SOD PRODUCTION****VENDETTA HERBICIDE DIRECTIONS****Seedling and Established Grasses**

			<b>APPLICATION TIMING AND SPECIFIC COMMENTS</b>	
<b>PRODUCT</b>	<b>RATE PER ACRE</b>	<b>RATE PER 1000 SQ. FT.</b>	<b>CROP</b>	<b>WEEDS</b>
VENDETTA Herbicide	1 to 2 pints	0.375 to 0.75 fl. oz.	Apply to established and newly seeded grasses grown for seed or sod production before the boot stage. Established grasses tolerant to Vendetta Herbicide include bentgrasses, Kentucky Bluegrass, Fescues, Ryegrass, Bermudagrass, St. Augustinegrass and Zoysiagrass. Vendetta Herbicide may also be used on seedling grasses such as Merion, Park, Delta, or common Kentucky Bluegrasses, Pennlawn, Chewings, Illahee or Alta Fescues, Orchard grass, Highland, Seaside or Astoria Bentgrasses, perennial Ryegrasses, Bahiagrass and Zoysiagrass.	Refer to the WEED LIST for a listing of susceptible broadleaf weeds.  Optimal control will be attained when weeds are treated in the seedling stage (less than 4 leaf stage, 2 inches in height, or 1 inch in diameter.)
VENDETTA Herbicide	Chemigation 2 pints/A only	0.75 fl. oz.	Apply to established and newly seeded grasses grown for seed or sod production before the boot stage. Apply through automated sprinkler irrigation systems with mechanical transfer loading system only. See MIXING, LOADING AND HANDLING INSTRUCTIONS section for complete details. Refer to the list of established grasses that are tolerant to Vendetta Herbicide.	

**RESTRICTIONS: Grasses grown for seed or sod production**

- Do not allow livestock to graze in treated areas or feed treated grasses to livestock.
- Do not apply Vendetta Herbicide to grasses grown for seed or sod production with backpack or hand-held application equipment.
- Do not apply more than 2 pints of Vendetta Herbicide (0.5 lb. MCPA acid equivalent) per acre in a single growing season.
- Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.
- Do not plant rotational crops within 30 days following application of Vendetta Herbicide.
- Do not apply more than 1.5 lbs. acid equivalent per acre per year of MCPA.

**FLAX (*Linum usitatissimum* only)****VENDETTA HERBICIDE DIRECTIONS**

<b>APPLICATION TIMING AND SPECIFIC COMMENTS</b>			
<b>PRODUCT</b>	<b>RATE</b>	<b>CROP</b>	<b>WEEDS</b>
VENDETTA Herbicide	0.5 to 0.9 pints/A	Apply to flax that is 2 to 8 inches in height. Do not apply Vendetta Herbicide to flax during or after the bud stage.	Apply to MOST SUSCEPTIBLE weeds that do not exceed the 4-leaf stage, 2 inches in height or 1 inch in diameter, whichever comes first.

**RESTRICTIONS and PRECAUTIONS: Flax (*Linum usitatissimum* only)**

- Do not apply if temperatures are expected to exceed 85° F at application or 3 days following application or crop injury may occur.
- Unacceptable crop injury may occur following Vendetta Herbicide application to flax grown on high organic, peat type soils.
- Application under high humidity conditions can injure flax.
- Unless otherwise instructed, do not apply Vendetta Herbicide to flax with crop oil concentrate, surfactants or nitrogen solutions.
- Do not use on ornamental flax.
- Do not apply more than 0.9 pints of Vendetta Herbicide (0.225 lb. MCPA acid equivalent) per acre in a single growing season.
- Do not plant rotational crops within 30 days following application of Vendetta Herbicide.
- Do not exceed 0.25 lb. acid equivalent per acre per year.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store at temperatures above 35° F. If allowed to freeze, remix before using.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Non-refillable containers (2.5, 30 & 55 gallons): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

**(Non-refillable <5 gallons): Triple rinse as follows:** Empty the remaining contents into application equipment or mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**(Non-refillable >5 gallons): Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable container (15 gallons, 30 gallons, 120 gallons and greater or bulk):** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

## Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

**ALL STATEMENTS MADE HEREIN ARE SUBJECT TO APPLICABLE LAW, AND TO THE EXTENT THERE IS ANY INCONSISTENCY OR CONTENTION, APPLICABLE LAW SHALL GOVERN.**

The Directions for Use of the product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

**Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS," AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE BEYOND WHAT IS STATED ON THE LABEL. Buyer and User accept all risks arising from any use of this product, including without limitation, uses contrary to label instructions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.**

Neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted, unless otherwise required by the law of the state of purchase, in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

WILBUR-ELLIS® Logo and VENDETTA® are registered trademarks of WILBUR-ELLIS Company.

WEEDAR and WEEDONE are registered trademarks of Nufarm Americas Inc.

Assert and Avenge are registered trademarks of BASF Corp.

Sencor is a registered trademark of Bayer AG, Germany.

Raindrop is a registered trademark of Delavan Corporation.

Ally, Express, Finesse, Glean, Harmony and Lexone are registered trademarks of E.I. du Pont de Nemours and Company.

Amber is a registered trademark of Syngenta Crop Protection.

Roundup is a registered trademark of Monsanto Company.

Curtail is a registered trademark of Dow AgroSciences LLC.

MANUFACTURED FOR:

**WILBUR-ELLIS COMPANY  
P.O. BOX 16458  
FRESNO, CA 93755**

F-031912

**NET CONTENTS:**